



 Medication: Clopidogrel (Plavix)
 PDN: 6982.02
 Last Updated: March 8, 2013
 PMD: Andrew Travers*
 PDC: Steven Carrigan*
 Page 1 of 2

CLOPIDOGREL (Plavix)

1.0 Classification

Platelet aggregation inhibitor

2.0 Mechanism of Action

 Irreversible blockade of adenosine diphosphate (ADP) receptor on platelet cell membranes, inhibiting platelet aggregation

3.0 Indications

Acute coronary syndrome (STEMI)

4.0 Contraindications

- Hypersensitivity
- Active pathological bleeding

5.0 Precautions

Administer with caution in patients on another platelet inhibitor

6.0 Route

May be given PO

7.0 Dosage

- As part of pre-hospital fibrinolysis: 75mg for patients 75 years old and above; 300mg for patients less than 75 years old
- For direct-to-PCI patients: 300mg

8.0 Supplied

• 75mg and 300mg tablets

9.0 May Be Given By

PCP/ICP/ACP/CCP

10.0 Adverse effects

- Bronchospasm
- Increased risk of bleeding
- Fever
- Neutropenia
- Thrombocytic Thrombocytopenic Purpura (TTP)

11.0 Special notes

- There is an increased risk of bleeding when clopidogrel is administered in conjunction with NSAIDs
- Pregnancy category C [if the patient will benefit from a Category C drug, it is generally used]

12.0 References

- Chest Pain Clinical Practice Guideline
- EHS STEMI Reperfusion Worksheet
- Compendium of Pharmaceuticals and Specialties (CPS)

*Electronically Signed Copyright © Emergency Health Services