

# RESPIRATORY WATCH

Week 44 (November 1 to November 7, 2015)\*

#### IN SUMMARY...

### **Activity levels**

 Central and eastern zones are reporting sporadic influenza activity. All other zones are reporting no activity.

# Laboratory-confirmed cases\*\*\*

- There have been 2 influenza cases this week, There have been 5 lab confirmed cases of influenza this season (1 influenza A H3 and 4 influenza A not subtyped).
- Positive test results were received for adenovirus, mycoplasma pneumonia, and pertussis.

# Severity


• There has been 1 ICU admission of laboratory confirmed influenza and 1 influenza deaths\*\*\* for the 2015-2016 influenza season.

# Syndromic surveillance

- The ILI rate for Nova Scotia this reporting period was 0.8.
- 100% of emergency departments reported ILI data for this period.

#### LABORATORY-CONFIRMED CASES

Figure 1: Number of respiratory specimens tested positive, and select percent positives, by report week, 2015-2016 season, Nova Scotia.



<sup>\*</sup>Data for this figure is obtained from provincial laboratories. All other data, unless otherwise stated, has been obtained from ANDS. Reporting lags may cause the data in this figure to not reconcile with others.

<sup>\*</sup>Reporting weeks run from Sunday to Saturday. The 2015-2016 influenza season is defined using PHAC's influenza surveillance weeks. This year runs from August 30, 2015 (Week 35) to August 27, 2016 (Week 34); \*\*Outbreak data is obtained from CNPHI, see appendix for definitions.; \*\*\* Deaths include individuals with a positive influenza test result, influenza may not have been the major contributing cause of death or hospitalization.

Figure 2: Number of reported lab-confirmed influenza cases by type and report week, 2015-2016 season, with trend-line comparison to 5-year average of all influenza cases, Nova Scotia.




Table 1: Number of total laboratory-confirmed influenza cases, current week and cumulative 2015-2016 season, by zones in Nova Scotia.

	Current Week					Cummulative 2015-2016				
Zone*		Influe	nza A		Influenza B	Influenza A				
	Total	A(H1) pdm09	A (H3)	A not subtyped		Total	A(H1) pdm09	A (H3)	A not subtyped	Influenza B
1 - Western	0	0	0	0	0	1	0	1	0	0
2 - Northern	0	0	0	0	0	0	0	0	0	0
3 - Eastern	1	0	0	1	0	1	0	0	1	0
4 - Central	1	0	0	1	0	3	0	0	3	0
Nova Scotia Total	2	0	0	2	0	5	0	1	1	0

<sup>\*</sup>Zones are defined in the appendix.

Table 2: Number of total laboratory-confirmed influenza cases, current week and cumulative 2015-2016 season, by age group in Nova Scotia.

	Current Week					Cummulative 2015-2016				
Zone*		Influer	nza A		Influenza B	Influenza A				
	Total	A(H1) pdm09	A (H3)	A not subtyped		Total	A(H1) pdm09	A (H3)	A not subtyped	Influenza B
0-4	0	0	0	0	0	0	0	0	0	0
5-19	0	0	0	0	0	0	0	0	0	0
20-44	0	0	0	0	0	0	0	0	0	0
45-64	1	0	0	1	0	2	0	0	2	0
65+	1	0	0	1	0	3	0	1	2	0
Nova Scotia Total	2	0	0	2	0	5	0	1	4	0




Figure 3: RSV cases by age group, cumulative 2015-2016 season, Nova Scotia.

# **SEVERITY**

Table 3: Hospitalizations, ICU admissions and deaths for influenza positive patients, current week and cumulative, 2015-2016 season, Nova Scotia.

Dothogon	Curre	ent surveillance	week	<b>Cumulative 2015-2016</b>		
Pathogen	Hospitalized*	ICU	Death**	Hospitalized	ICU	Death
Influenza A (H1) pdm09	0	0	0	0	0	0
Influenza A unsubtyped	0	0	1	0	1	1
Influenza A (H3)	0	0	0	1	0	0
Influenza B	0	0	0	0	0	0
Nova Scotia Total	0	0	1	1	1	1

<sup>\*</sup>Hospitalizations do not include ICU admissions; \*\*Deaths include individuals with a positive influenza test result, influenza may not have been the major contributing cause of death or hospitalization.\*

#### **SYNDROMIC SURVEILLANCE**

Figure 4: Percentage of emergency room visits due to ILI by week, 2015-2016 season, with trend-line comparison to 2014-2015, Nova Scotia.




Table 4: Current week ILI reporting from emergency departments by zones, 2015-2016 season, Nova Scotia.

Zone	ILI (%)	Number of reporting ERs	Number of ERs	ERs reporting (%)	
1 - Western	0.6	9	9	100.0	
2 - Northern	0.2	8	8	100.0	
3 - Eastern	0.9	14	14	100.0	
4 - Central	1.3	7	7	100.0	
IWK	5.9	2	2	100.0	
Nova Scotia Total (excl IWK)	0.5	38	38	100.0	
Nova Scotia Total (incl IWK)	0.8	40	40	100.0	

# OTHER RESPIRATORY PATHOGENS

Table 5: Total number of specimens tested and number (%) positive for other respiratory pathogens, by report week and cumulative season, Nova Scotia, 2015–2016

Pathogen		Surveillance	Week	Cumulative Season-to-Date Totals			
. aanogon	n tested	n positive	% positive	n tested	n positive	% positive	
Adenovirus	6	1	16.7	176	6	3.4	
Bocavirus	6	0	0.0	176	0	0.0	
Chlamydophila pneumoniae	35	0	0.0	161	1	0.6	
Coronavirus	6	0	0.0	176	0	0.0	
Enterovirus	6	0	0.0	176	1	0.6	
Metapneumovirus	6	0	0.0	176	0	0.0	
Mycoplasma pneumoniae	35	12	34.3	161	48	29.8	
Parainfluenza	6	0	0.0	176	4	2.3	
Pertussis	27	1	3.7	245	29	11.8	
Respiratory Syncytial Virus	6	0	0.0	276	0	0.0	
Rhinovirus	6	0	0.0	176	32	18.2	

# APPENDIX: DEFINITIONS USED IN INFLUENZA SURVEILLANCE, AND USEFUL LINKS, 2015-2016

#### **ACRONYM LIST**

**CNPHI** Canadian Network for Public Health Intelligence

DHA District Health Authority
 ICU Intensive care unit
 ILI Influenza-like illness
 LTCF Long-term care facility
 RSV Respiratory syncytial virus

# **ILI CASE DEFINITION**

Acute onset of respiratory illness with fever and cough and with one or more of the following – sore throat, arthralgia, myalgia or prostration which is likely due to influenza. In children under 5, gastrointestinal symptoms may also be present. In patients under 5 or 65 and older, fever may not be prominent.

#### **OUTBREAK DEFINITIONS**

Schools and daycares	Greater than 10% absenteeism (or absenteeism that is higher (e.g. >5-10%) than expected level as determined by the school or public health authority) which is likely
	due to ILI.
Hospitals and	Two or more cases of ILI within a seven-day period, including at least one laboratory
residential institutions	confirmed case. Institutional outbreaks should be reported within 24 hours of
	identification. Residential institutions include, but are not limited to, long-term care
	facilities (LTCF) and prisons.
Other settings	Two or more cases of ILI within a seven-day period, including at least one laboratory
	confirmed case (i.e. closed communities).

#### NATIONAL FLUWATCH DEFINITIONS FOR INFLUENZA ACTIVITY LEVELS

No activity	No laboratory-confirmed influenza detections in the reporting week, however, sporadically occurring ILI* may be reported					
Sporadic	Sporadically occurring ILI* and lab confirmed influenza detection(s) with <b>no outbreaks</b> detected within the influenza surveillance region					
Localized	<ol> <li>Evidence of increased ILI* and</li> <li>lab confirmed influenza detection(s) together with</li> <li>outbreaks occurring in schools, hospitals, residential institutions and/or other types of facilities occurring in less than 50% of the influenza surveillance region</li> </ol>					
Widespread	<ul> <li>(1) Evidence of increased ILI* and</li> <li>(2) lab confirmed influenza detection(s) together with</li> <li>(3) outbreaks occurring in schools, hospitals, residential institutions and/or other types of facilities occurring in greater than or equal to 50% of the influenza surveillance region</li> </ul>					

#### LINKS TO OTHER WEEKLY INFLUENZA REPORTING BODIES

Canada: http://www.phac-aspc.gc.ca/fluwatch/

World: https://www.who.int/influenza/surveillance\_monitoring/updates/latest\_update\_GIP\_surveillance/en/index.html

US: www.cdc.gov/flu/weekly