

# RESPIRATORY WATCH

Week 3 (January 17 to January 23, 2016)\*

### IN SUMMARY...

### **Activity levels\*\***

Central (Zone 4) is reporting localized activity.

# Laboratory-confirmed cases\*\*\*

- There were 2 influenza case reported this week. There have been 13 lab confirmed cases of influenza this season (1 influenza A H3 and 12 influenza A not subtyped).
- Positive test results were received for adenovirus, mycoplasma pneumonia, pertussis and RSV.

### Severity


 There has been 1 ICU admission of laboratory confirmed influenza and 1 influenza death\*\*\*\* for the 2015-2016 influenza season.

# Syndromic surveillance

- The ILI rate for Nova Scotia this reporting period was 0.9.
- 100% of emergency departments reported ILI data for this period.

#### LABORATORY-CONFIRMED CASES

Figure 1: Number of respiratory specimens tested positive, and select percent positives, by report week, 2015-2016 season. Nova Scotia.



<sup>\*</sup>Data for this figure is obtained from provincial laboratories. All other data, unless otherwise stated, has been obtained from ANDS. Reporting lags may cause the data in this figure to not reconcile with others.

<sup>\*</sup>Reporting weeks run from Sunday to Saturday. The 2015-2016 influenza season is defined using PHAC's influenza surveillance weeks. This year runs from August 30, 2015 (Week 35) to August 27, 2016 (Week 34); \*\*Activity level data is obtained from CNPHI, see appendix for definitions.; \*\*\*Week 51 does not include laboratory data from the IWK \*\*\*\*\*Deaths include individuals with a positive influenza test result, influenza may not have been the major contributing cause of death or hospitalization.

Figure 2: Number of reported lab-confirmed influenza cases by type and report week, 2015-2016 season, with trend-line comparison to 5-year average of all influenza cases, Nova Scotia.




Table 1: Number of total laboratory-confirmed influenza cases, current week and cumulative 2015-2016 season, by zones in Nova Scotia.

	Current Week				Cummulative 2015-2016					
Zone* Influenza A					Influenza A					
	Total	A(H1) pdm09	A (H3)	A not subtyped	Influenza B	Total	A(H1) pdm09	A (H3)	A not subtyped	Influenza B
1 - Western	1	0	1	0	0	1	0	1	0	0
2 - Northern	1	0	0	1	0	1	0	0	1	0
3 - Eastern	1	0	0	1	0	1	0	0	1	0
4 - Central	10	0	0	10	0	10	0	0	10	0
Nova Scotia Total	13	0	1	12	0	13	0	1	12	0

<sup>\*</sup>Zones are defined in the appendix.

Table 2: Number of total laboratory-confirmed influenza cases, current week and cumulative 2015-2016 season, by age group in Nova Scotia.

	Current Week					Cummulative 2015-2016				
Age		Influer	nza A		Influenza B	Influenza A				
3.	Total	A(H1) pdm09	A (H3)	A not subtyped		Total	A(H1) pdm09	A (H3)	A not subtyped	Influenza B
0-4	0	0	0	0	0	2	0	0	2	0
5-19	0	0	0	1	0	3	0	0	3	0
20-44	0	0	0	0	0	0	0	0	0	0
45-64	1	0	0	1	0	5	0	0	5	0
65+	0	0	0	0	0	3	0	1	2	0
Nova Scotia Total	2	0	0	2	0	13	0	1	12	0




Figure 3: RSV cases by age group, cumulative 2015-2016 season, Nova Scotia.

## **SEVERITY**

Table 3: Hospitalizations, ICU admissions and deaths for influenza positive patients, current week and cumulative, 2015-2016 season, Nova Scotia.

Dothoron	Curre	nt surveillance	week	Cumulative 2015-2016			
Pathogen	Hospitalized*	ICU	Death**	Hospitalized	ICU	Death	
Influenza A (H1) pdm09	0	0	0	0	0	0	
Influenza A unsubtyped	2	0	0	9	1	1	
Influenza A (H3)	0	0	0	1	0	0	
Influenza B	0	0	0	0	0	0	
Nova Scotia Total	2	0	0	10	1	1	

<sup>\*</sup>Hospitalizations do not include ICU admissions; \*\*Deaths include individuals with a positive influenza test result, influenza may not have been the major contributing cause of death or hospitalization.\*

#### **SYNDROMIC SURVEILLANCE**

Figure 4: Percentage of emergency room visits due to ILI by week, 2015-2016 season, with trend-line comparison to 2014-2015, Nova Scotia.




Table 4: Current week ILI reporting from emergency departments by zones, 2015-2016 season, Nova Scotia.

Zone	ILI (%)	Number of reporting ERs Number of ERs		ERs reporting (%)
1 - Western	0.4	9	9	100.0
2 - Northern	0.2	8	8	100.0
3 - Eastern	0.7	14	14	100.0
4 - Central	1.6	7	7	100.0
IWK	6.6	2	2	100.0
Nova Scotia Total (excl IWK)	0.4	38	38	100.0
Nova Scotia Total (incl IWK)	0.9	40	40	100.0

# OTHER RESPIRATORY PATHOGENS

Table 5: Total number of specimens tested and number (%) positive for other respiratory pathogens, by report week and cumulative season, Nova Scotia, 2015–2016

Pathogen		Surveillance \	Week	Cumulative Season-to-Date Totals			
ratiogen	n tested	n positive	% positive	n tested	n positive	% positive	
Adenovirus	6	1	16.7	266	12	4.5	
Bocavirus	6	0	0.0	266	3	1.1	
Chlamydophila pneumoniae	15	0	0.0	473	2	0.4	
Coronavirus	6	0	0.0	266	0	0.0	
Enterovirus	6	0	0.0	266	1	0.4	
Metapneumovirus	6	0	0.0	266	0	0.0	
Mycoplasma pneumoniae	15	3	20.0	473	122	25.8	
Parainfluenza	6	0	0.0	266	10	3.8	
Pertussis	16	2	12.5	491	43	8.8	
Respiratory Syncytial Virus	39	4	10.3	774	7	0.9	
Rhinovirus	6	0	0.0	266	44	16.5	

### APPENDIX: DEFINITIONS USED IN INFLUENZA SURVEILLANCE, AND USEFUL LINKS, 2015-2016

#### **ACRONYM LIST**

CNPHI Canadian Network for Public Health Intelligence

DHA District Health Authority
 ICU Intensive care unit
 ILI Influenza-like illness
 LTCF Long-term care facility
 RSV Respiratory syncytial virus

#### **ILI CASE DEFINITION**

Acute onset of respiratory illness with fever and cough and with one or more of the following – sore throat, arthralgia, myalgia or prostration which is likely due to influenza. In children under 5, gastrointestinal symptoms may also be present. In patients under 5 or 65 and older, fever may not be prominent.

#### **OUTBREAK DEFINITIONS**

Schools and daycares	Greater than 10% absenteeism (or absenteeism that is higher (e.g. >5-10%) than expected level as determined by the school or public health authority) which is likely due to ILI.
Hospitals and residential institutions	Two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case. Institutional outbreaks should be reported within 24 hours of identification. Residential institutions include, but are not limited to, long-term care facilities (LTCF) and prisons.
Other settings	Two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case (i.e. closed communities).

#### NATIONAL FLUWATCH DEFINITIONS FOR INFLUENZA ACTIVITY LEVELS

No activity	No laboratory-confirmed influenza detections in the reporting week, however,
	sporadically occurring ILI* may be reported
Sporadic	Sporadically occurring ILI* and lab confirmed influenza detection(s) with <b>no</b>
	outbreaks detected within the influenza surveillance region
Localized	<ul> <li>(1) Evidence of increased ILI* and</li> <li>(2) lab confirmed influenza detection(s) together with</li> <li>(3) outbreaks occurring in schools, hospitals, residential institutions and/or other types of facilities occurring in less than 50% of the influenza surveillance region</li> </ul>
Widespread	<ul> <li>(1) Evidence of increased ILI* and</li> <li>(2) lab confirmed influenza detection(s) together with</li> <li>(3) outbreaks occurring in schools, hospitals, residential institutions and/or other types of facilities occurring in greater than or equal to 50% of the influenza surveillance region</li> </ul>

#### LINKS TO OTHER WEEKLY INFLUENZA REPORTING BODIES

Canada: <a href="http://www.phac-aspc.gc.ca/fluwatch/">http://www.phac-aspc.gc.ca/fluwatch/</a>

World: https://www.who.int/influenza/surveillance monitoring/updates/latest update GIP surveillance/en/index.html

US: www.cdc.gov/flu/weekly