


Summary of Nova Scotia surveillance findings, for the period ending January 19, 2013:

- Twenty four influenza A positive lab results were reported this week.
- Other respiratory pathogen activity continues. Positive results were received for coronavirus, metapneumovirus, mycoplasma pneumoniae, rhinovirus and RSV.
- The ILI rate for Nova Scotia for this reporting period was 1.4. Eighty-two percent of ER sites reported ILI data this week.
- Three specimens were submitted through the sentinel swabbing program. Two from DHA 7 and 1 from DHA 8.
- Sentinel physician data was not received at time of report production.

Figure 1: Summary of laboratory detected circulating respiratory pathogens, Nova Scotia, 2012–2013



See Table 4 for further details regarding other respiratory pathogens.




RESPIRATORY WATCH

Week 3 (January 13 to January 19, 2013)

INFLUENZA AND INFLUENZA-LIKE ILLNESS

Figure 2: Influenza and ILI activity, surveillance week 3, Nova Scotia



RESPIRATORY WATCH

Week 3 (January 13 to January 19, 2013)

Figure 3: Number of reported lab-confirmed influenza cases by type and report week, Nova Scotia, 2012–2013





Figure 4: Number of specimens tested for influenza and percent positive, Nova Scotia Provincial Public Health Laboratory Network, 2012–2013*



*Data presented in this figure refers to week specimen was tested.

Table 1: Influenza case counts by DHA, current surveillance week and cumulative, Nova Scotia, 2012–2013

	DHA 1	DHA 2	DHA 3	DHA 4	DHA 5	DHA 6	DHA 7	DHA 8	DHA 9	Nova Scotia
Influenza A 2009 pH1N1										
Current Week	0	0	0	0	0	0	0	0	0	0
Cumulative 2012 - 2013	0	0	0	0	0	0	0	0	0	0
Influenza A (not yet sub-typed)										
Current Week	2	0	0	1	0	2	0	2	8	15
Cumulative 2012 - 2013	4	3	3	1	4	5	4	4	28	56
Influenza A Seasonal (H3)										
Current Week	0	0	0	0	0	0	0	0	0	0
Cumulative 2012 - 2013	0	0	0	0	0	0	0	0	0	0
Influenza B										
Current Week	0	0	0	0	0	0	0	0	0	0
Cumulative 2012 - 2013	0	0	0	0	0	0	0	0	2	2

RESPIRATORY WATCH

Week 3 (January 13 to January 19, 2013)

Figure 5: Influenza rate per 100,000 population by type and age group, cumulative, Nova Scotia, 2012–2013




Figure 6: Influenza rate per 100,000 population by type and DHA, cumulative, Nova Scotia, 2012–2013




Table 2: ILI reporting from emergency departments and FluWatch sentinel physicians, and Sentinel Swabbing Specimen Submissions, Nova Scotia, 2012-2013

	ER SURVEILLANCE		SENTINEL SURVEILLANCE*		SENTINEL SWABBING	
	%ILI	Reporting ERs	%ILI	Reporting Sentinels	# Swabs	Sites Submitting Specimens
DHA 1	0.9	3 of 3	–	0 of 6	0	0 of 1
DHA 2	0.0	3 of 3	–	0 of 0	0	0 of 1
DHA 3	0.9	3 of 5	–	0 of 1	0	0 of 2
DHA 4	1.1	2 of 2	–	0 of 0	0	0 of 2
DHA 5	1.7	5 of 5	–	0 of 2	0	0 of 1
DHA 6	1.1	1 of 1	–	0 of 2	0	0 of 1
DHA 7	2.4	6 of 6	–	0 of 1	2	1 of 2
DHA 8	2.1	6 of 8	–	0 of 4	1	1 of 3
DHA 9	0.6	4 of 7	–	0 of 14		
IWK	5.6	1 of 1				
Nova Scotia (excl. IWK)†	1.1	33 of 40	82.5%		3	2 of 12
Nova Scotia (incl. IWK)	1.4	34 of 41	82.9%	0 of 30	0.0%	


*Flu watch sentinels

†Excludes the children's ER from IWK

RESPIRATORY WATCH

Week 3 (January 13 to January 19, 2013)

Figure 7: Percentage of ER visits with ILI, Nova Scotia, 2012–2013



RESPIRATORY WATCH

Week 3 (January 13 to January 19, 2013)

RESPIRATORY SYNCYTIAL VIRUS (RSV)

Figure 8: Number of positive RSV specimens by report week, Nova Scotia, 2012–2013





Figure 9: Cumulative number of positive RSV specimens by age group and sex, Nova Scotia, 2012-2013



RESPIRATORY WATCH

Week 3 (January 13 to January 19, 2013)

OTHER RESPIRATORY PATHOGENS

Table 3: Total number of specimens tested and number (%) positive for other respiratory pathogens, by report week and cumulative season, Nova Scotia, 2012–2013

Number and percent positive for:	Surveillance Week			Cumulative		
	n tested	n positive	% positive	Season-to-Date	Totals	
	n tested	n positive	% positive	n tested	n positive	% positive
Adenovirus	23	0	0.0	329	0	0.0
Bocavirus	23	0	0.0	329	1	0.3
Chlamydomphila pneumoniae	15	0	0.0	277	23	8.3
Coronavirus	23	3	13.0	329	8	2.4
Enterovirus	23	0	0.0	329	4	1.2
Metapneumovirus	23	1	4.3	329	5	1.5
Mycoplasma pneumoniae	15	3	20.0	277	60	21.7
Parainfluenza	23	0	0.0	329	15	4.6
Pertussis	12	0	0.0	161	14	8.7
Respiratory syncytial virus A	0	0	0.0	308	8	2.6
Respiratory syncytial virus B	0	0	0.0	308	0	0.0
Respiratory syncytial virus not typed	96	7	7.3	245	16	6.5
Rhinovirus	23	4	17.4	329	47	14.3

RESPIRATORY WATCH

Week 3 (January 13 to January 19, 2013)

APPENDIX: Definitions used in Influenza Surveillance, 2012-2013

1) ILI in the general population:

Acute onset of respiratory illness with fever and cough and with one or more of the following - sore throat, arthralgia, myalgia, or prostration which is likely due to influenza. In children under 5, gastrointestinal symptoms may also be present. In patients under 5 or 65 and older, fever may not be prominent.

2) Outbreaks of influenza / ILI by setting:

Schools and Daycares:

Greater than 10% absenteeism (or absenteeism that is higher (e.g. >5-10%) than expected level as determined by school or public health authority) which is likely due to ILI.

Hospitals and residential institutions:

Two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case. Institutional outbreaks should be reported within 24 hours of identification. Residential institutions include but not limited to long-term care facilities (LTCF) and prisons.

Other Settings:

Two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case; i.e. closed communities.

3) National FluWatch Definitions for Influenza Activity Levels:

Influenza activity levels are defined as:

- | | |
|-------------------------|--|
| 1 = No activity: | i.e. no laboratory-confirmed influenza detections in the reporting week, however, sporadically occurring ILI* may be reported |
| 2 = Sporadic: | sporadically occurring ILI* and lab confirmed influenza detection(s) with no outbreaks detected within the influenza surveillance region† |
| 3 = Localized: | (1) evidence of increased ILI* and
(2) lab confirmed influenza detection(s) together with
(3) outbreaks in schools, hospitals, residential institutions and/or other types of facilities occurring in less than 50% of the influenza surveillance region † |
| 4 = Widespread: | (1) evidence of increased ILI* and
(2) lab confirmed influenza detection(s) together with
(3) outbreaks in schools, hospitals, residential institutions and/or other types of facilities occurring in greater than or equal to 50% of the influenza surveillance region † |

* ILI data may be reported through sentinel physicians, emergency room visits or health line telephone calls.

† Sub-regions within the province or territory as defined by the provincial/territorial epidemiologist.

RESPIRATORY WATCH

Week 3 (January 13 to January 19, 2013)

- 4) District Health Authorities (DHAs), Nova Scotia:
 - DHA 1 – South Shore Health
 - DHA 2 – South West Health
 - DHA 3 – Annapolis Valley Health
 - DHA 4 – Colchester East Hants Health Authority
 - DHA 5 – Cumberland Health Authority
 - DHA 6 – Pictou County Health Authority
 - DHA 7 – Guysborough Antigonish Strait Health Authority
 - DHA 8 – Cape Breton District Health Authority
 - DHA 9 – Capital Health