

# PARATYPHOID FEVER

## Case definition

### CONFIRMED CASE

Laboratory confirmation of infection with or without clinical illness:

- isolation of *Salmonella paratyphi* from an appropriate clinical specimen (i.e., sterile site, deep tissue wounds, stool, vomit or urine).

### PROBABLE CASE

Clinical illness in a person who is epidemiologically linked to a confirmed case.

## Causative agent

*Salmonella paratyphi*

## Source

Stool and/or urine of infected person. Rarely, domestic animals are a reservoir. A small number of people may become permanent carriers.

## Incubation

Dependent on infectious dose, generally from 1 day to 10 days.

## Transmission

- Fecal-oral from person-to-person or by ingestion of food or water contaminated by feces or urine of the infected person or domestic animal.
- Consumption of shellfish harvested from sewage contaminated waters or consumption of uncooked fruits and vegetables fertilized with human waste. Flies may be a vehicle for the contamination of food.

## Communicability

Usually from the first week of illness throughout convalescence, commonly 1-2 weeks. Carrier state is possible; short term fecal carriage is more likely than chronic carrier state. Chronic carrier state (< 5% of population) is usually linked to the biliary or urinary tract and should be distinguished from short-term fecal carriage.

## Symptoms

Characterized by insidious onset of sustained fever, headache, malaise, anorexia, splenomegaly, diarrhea (more common in children) or constipation (more common

in adults] and non-productive cough. Relative bradycardia and occasionally a transient, macular rash of rose-coloured spots can be seen on the trunk.

Atypical presentations occur, and the severity of illness varies, with a generally milder presentation than *S. typhi* and a much lower case fatality rate.

## Diagnostic testing

Stool, urine, bone marrow or blood for culture. Organisms are often absent from stool.

Testing may also need to be completed for schistosomiasis if case history indicates travel or having lived in an endemic area. As schistosome infections can be hepatic, intestinal and/or urinary, multiple sample types may be required for definitive diagnosis.

## Treatment

Treatment may include antibiotics and/or corticosteroids.

# PUBLIC HEALTH MANAGEMENT & RESPONSE

---

## Case management

Follow up the case using the following steps:

1. Contact the primary care provider to obtain clinical information on the case.
2. Interview the case, review clinical information, determine food history, travel history and travel activities, employment, potential source of exposure and determine any contacts that may require investigation [see "[Contact tracing](#)" section].
3. Educate the case and/or family about Paratyphoid Fever and prevention measures, providing access to website, general information, etc.
4. Implement the necessary exclusions as per the "[Exclusion of cases and carriers](#)" section for those cases identifying as belonging to one or more risk group[s]. For cases that are not listed in either of the risk groups, recommend that the case remain at home until 48 hours after stools have returned to normal and 48 hours after stopping the use of anti-diarrheal medication.
5. If the case has no travel history and identifies consuming shellfish, especially shellfish harvested from an area possibly contaminated with sewage, or raw fruits and vegetables purchased at a food establishment, contact a Food Safety Specialist with the Department of Environment.
6. Document the information on the Enteric Case Report Form.

## Exclusion of cases and carriers

Individuals who continue to shed *Salmonella paratyphi* for one year or more are considered to be chronic carriers.

Case management of carriers employed in any of the below high-risk groups should be done in consultation with the regional MOH as redeployment of staff to lower risk activities may be possible.

Exclude cases and carriers in the risk groups below:

Risk Group	Criteria for Exclusion
Food handlers	Until 3 negative stool samples have been obtained at least 24 hours apart AND at least 48 hours after discontinuance of antibiotics AND one negative urine culture if travelled or lived in schistosomiasis endemic area.
Health care, child care or other staff who have contact with susceptible persons	Until 3 negative stool samples have been obtained at least 24 hours apart AND at least 48 hours after discontinuance of antibiotics AND one negative urine culture if travelled or lived in schistosomiasis endemic area.
Children attending child care	Until 3 negative stool samples have been obtained at least 24 hours apart AND at least 48 hours after discontinuance of antibiotics AND one negative urine culture if travelled or lived in schistosomiasis endemic area.
Carrier(s) [both symptomatic and asymptomatic] employed in: <ul style="list-style-type: none"><li>• food handling</li><li>• child care*</li><li>• health care and/or other staff who have contact with susceptible persons</li></ul> * Inclusive of those attending child care.	Until 3 negative stool samples have been obtained at least 24 hours apart AND at least 48 hours after discontinuance of antibiotics AND one negative urine culture if travelled or lived in schistosomiasis endemic area.  If any of these samples are positive, repeat cultures at weekly intervals for 8 weeks until 3 consecutive samples are negative.  If 3 consecutive negative samples are not obtained after 8 weekly samples, repeat cultures monthly for up to 10 months until 3 consecutive samples are negative.  If 3 consecutive negative samples are not obtained after 10 monthly samples the person is considered a chronic carrier.

**Note:** Ensure that all samples submitted to the laboratory for testing are labelled "Public Health management requirement to inform exclusion".

## Education of cases and carriers

Offer the following information:

- Ensure cases belonging to a high-risk group are aware of exclusion criteria.
- Remind cases about the importance of hand hygiene in stopping the spread of paratyphoid fever and to wash hands before preparing food and after using the bathroom and changing diapers.
- Inform the case about the potential to infect contacts and provide information on how to minimize transmission to others; including household and close contacts, including sexual contacts.
- Recommend that cases infected with paratyphoid fever or any other gastrointestinal illness should not prepare or serve food to other people [for food handlers see “[Exclusion of cases and carriers](#)” section].

See the [General Information Sheet](#) for further information on preventing the transmission of paratyphoid fever.

## Contact tracing

Contact tracing should be initiated as part of case management if symptomatic contacts or contacts that belong to any of the risk groups identified in the “[Exclusion of contacts](#)” section are identified by the case.

### Definition of a contact

A contact is a person who has had exposure to a case during the period of communicability and is at risk of infection by the fecal-oral route by either person-to-person contact or the ingestion of contaminated food or water.

Contacts include:

- Household contacts (those living in the same residence)
- Close contacts including sexual contacts and persons who may have had hand-to-mouth contact with the case, such as sharing meals the case has prepared.
- All members of a travel group associated with a case [e.g., those who travelled together to the same location(s), not just on the same flight]

### Exclusion of contacts

Close contacts (household and sexual) of cases and carriers not employed in any of the risk groups listed below should be provided information about disease transmission and appropriate infection prevention and control measures, including seeking prompt medical assessment and notifying Public Health if they become symptomatic.

Exclude contacts in the risk groups below:

Risk Group	Criteria for Exclusion
<p>Co-traveller(s) (both symptomatic and asymptomatic) employed in:</p> <ul style="list-style-type: none"> <li>• food handling</li> <li>• child care*</li> <li>• health care and/or other staff who have contact with susceptible persons</li> </ul> <p>* Inclusive of those attending child care.</p>	<p>Until 2 negative stool samples have been obtained at least 24 hours apart AND at least 48 hours after discontinuance of antibiotics AND one negative urine culture if travelled or lived in schistosomiasis endemic area.</p> <p>Note: If any of the culture specimens are positive for <i>Salmonella paratyphi</i> then treat as a case.</p>
<p>Symptomatic close contacts of cases and carriers (household and sexual) employed in:</p> <ul style="list-style-type: none"> <li>• food handling</li> <li>• child care*</li> <li>• health care and/or other staff who have contact with susceptible persons</li> </ul> <p>* Inclusive of those attending child care.</p>	<p>Until 2 negative stool samples have been obtained at least 24 hours apart AND at least 48 hours after discontinuance of antibiotics AND one negative urine culture if travelled or lived in schistosomiasis endemic area.</p> <p>Note: If any of the culture specimens are positive for <i>Salmonella paratyphi</i> then treat as a case.</p>
<p>Asymptomatic close contacts of cases (household and sexual) employed in:</p> <ul style="list-style-type: none"> <li>• food handling</li> <li>• child care*</li> <li>• health care and/or other staff who have contact with susceptible persons</li> </ul> <p>* Inclusive of those attending child care.</p>	<p>Collect one screening stool sample.</p> <p>Exclusion not necessary while awaiting culture results.</p> <p>Note: If any of the culture specimens are positive for <i>Salmonella paratyphi</i> then treat as a case.</p>

**[continued on next page]**

Risk Group	Criteria for Exclusion
<p>Asymptomatic close contacts (household and sexual) of carriers employed in:</p> <ul style="list-style-type: none"> <li>• food handling,</li> <li>• child care*</li> <li>• health care and/or other staff who have contact with susceptible persons</li> </ul> <p>* Inclusive of those attending child care.</p>	<p>Are not excluded and no stool specimens are required, however are advised to seek prompt medical assessment and notify Public Health if they become symptomatic.</p> <p>Close contacts of carriers should be provided with information about symptoms, disease transmission, appropriate infection prevention and control measures and immunization (where applicable).</p>

Note: Ensure that all samples submitted to the laboratory for testing are labelled “Public Health management requirement to inform exclusion”.

### Education of contacts

If Public Health is notifying contacts, inform the contacts of the following:

- Their potential exposure
- An explanation of the illness (description of the disease, symptoms, etc.)
- The range of clinical presentation
- Incubation period
- Report to Public Health if they become symptomatic.

See the [General Information Sheet](#) for further information on preventing the transmission of paratyphoid fever.

### Outbreak control

Consult the [Outbreak Response Plan](#) for further guidance if an outbreak is suspected.

For outbreaks in child care settings also refer to the [Guidelines for Communicable Disease Prevention and Control for Child Care Settings](#).

For outbreaks in Long-Term Care Facilities also refer to Infection Prevention and Control Nova Scotia’s (IPCNS) [Infection Prevention and Control: Guidelines for Long-Term Care Facilities](#).

## Surveillance forms

[novascotia.ca/dhw/populationhealth/surveillanceguidelines/NS\\_Notifiable\\_Disease\\_Surveillance\\_Case\\_Report\\_Form.docx](http://novascotia.ca/dhw/populationhealth/surveillanceguidelines/NS_Notifiable_Disease_Surveillance_Case_Report_Form.docx)

[novascotia.ca/dhw/populationhealth/surveillanceguidelines/Enteric\\_Case\\_Report\\_Form.pdf](http://novascotia.ca/dhw/populationhealth/surveillanceguidelines/Enteric_Case_Report_Form.pdf)

## General Information Sheet

## References

Alberta Health. April 2014. Public Health Notifiable Disease Management Guidelines - Paratyphoid Fever. [health.alberta.ca/documents/Guidelines-Paratyphoid-Fever-2014.pdf](http://health.alberta.ca/documents/Guidelines-Paratyphoid-Fever-2014.pdf)

BC Centre for Disease Control. Exclusion of Enteric Cases and their Contacts from High Risk Settings. May 2013 [bccdc.ca/NR/rdonlyres/56C97580-5A9C-41C5-8F22-3818337C55A5/0/EntericCasesandtheirContacts\\_May2013.pdf](http://bccdc.ca/NR/rdonlyres/56C97580-5A9C-41C5-8F22-3818337C55A5/0/EntericCasesandtheirContacts_May2013.pdf)

Centers for Disease Control and Prevention. Schistosomiasis Endemic Area map. [cdc.gov/travel-static/yellowbook/2016/map\\_3-12.pdf](http://cdc.gov/travel-static/yellowbook/2016/map_3-12.pdf)

*Control of Communicable Diseases Manual*, 20th edition. 2015. David Heymann, MD, editor.

*Provincial Microbiology User's Manual*. [cdha.nshealth.ca/pathology-laboratory-medicine](http://cdha.nshealth.ca/pathology-laboratory-medicine)

Public Health Agency of Canada. [2009]. Case Definitions for Communicable Diseases under National Surveillance.

*Red Book. 2012 Report of the Committee on Infectious Diseases*, 29th edition. American Academy of Pediatrics.