


# Geological Map for Part of NTS 11F/05, Guysborough Area, Nova Scotia

J. B. Murphy and R. J. Rice

Scale 1:50 000



Halifax, Nova Scotia



© Her Majesty the Queen in Right of the Province of Nova Scotia 2005

## Selected References

Benson, D.G. 1967. Geology of the Hopewell map area, Nova Scotia; Geological Survey of Canada, Memoir 343, 58 p.  
Benson, D.G. 1974. Geology of the Antigonish Highlands, Nova Scotia; Geological Survey of Canada, Memoir 376, 92 p.

## Descriptive Text

The St. Marys Basin, central mainland Nova Scotia, is dominated by a Late Devonian(?) - Early Carboniferous intracratonic aluvial fan-fluvial lacustrine basin-fill sequence that occupies the current boundary between the Meguma and Avalon terranes of the Canadian Appalachians. The Basin rocks belong to the Horton Group which is divided into six, largely laterally equivalent, formations. The stratigraphically lowest rocks are predominantly exposed in the central part of the Basin in a series of an echelon anticlinal closures (11E/06-08).

In the northwestern part of this map (11F/05), the Lochiel Formation typically consists of sandstones which are grey or green, fine- to very coarse-grained, micaceous and feldspathic, interbedded with minor granitoides and dark grey siltstone. The sandstones are typically 1 to 2 m in thickness and abundant fining-upward cycles have bases defined by granitoides beds. Sandstone beds occur in measured intervals up to 17 m in thickness. Parallel lamination, ripple crosslamination and large-scale crossbedding are common in the sandstones. The crossbeds are both trough (dominant) and planar with set thicknesses varying from 0.1 to 0.6 m. Cores of trough cross-stratification can be up to several metres thick. Grey weathering siltstone, similar to that in the Barrens Hills Formation, is sparse and occurs in massive, highly fractured intervals 1-2 m thick. Parallel lamination and ripple crosslamination are present in the siltstone, but may be partially masked by fracturing. Towards the southern and eastern flanks of the Basin (this map), the sequence is dominated by beds that coarsen and thicken upward and were predominantly deposited in alluvial fans (Cross Brook and West River St. Marys formations) derived from the Meguma Terrane to the south.

The Cross Brook Formation overlies the Lochiel Formation and consists of grey-green weathering sandstone interstratified with less abundant grey-green weathering siltstone, shale, conglomerate and rare limestone. The Formation can be distinguished from the Lochiel Formation by the abundance of green and red metamorphic clasts that are clearly derived from the Meguma Terrane to the south. The sandstones are typically fine- to coarse-grained, micaceous, variably feldspathic, and locally contain varying amounts of granule-sized quartz and sedimentary/metasedimentary lithic fragments. They generally are thinly bedded and commonly display ripple crosslamination, and trough cross-stratification in sets up to 1 m thick. Pebble conglomerates are polygenic, poorly sorted, and framework supported. Dominant clast types in the sandstones and conglomerates include psammite, pelite, micaceous granite, vein quartz and muscovite. Less abundant, sedimentary clasts include laminated, carbonaceous mudstone (possibly of algal origin), siltstone, sandstone, carbonate and organic debris, all of which are interpreted as intra-basin detritus. Most clasts are subangular to subrounded. Siltstone and shale beds are grey weathering, micaceous and display parallel lamination and ripple crosslamination.

The West River St. Marys Formation consists of reddish-brown to grey-brown weathering conglomerate interstratified with grey-brown weathering sandstone. The Formation outcrops along the southern flank of the central and eastern portions of the Basin. The red-brown to grey-brown weathering conglomerate contains clasts ranging from pebbles to boulders. The rock is dominantly framework- to locally matrix-supported with a roughly 80:20 framework to matrix ratio. It is very poorly to poorly sorted, may contain imbricated pebbles, or display crude inverse- to normal-grading. The conglomerate contains numerous major scour surfaces and fines upward to large pebble-size in the top of the measured section. Basal contacts with sandstone lenses are erosional. Clast shape and size vary with composition. Sedimentary clasts are largely sandstone to 20 cm, shale/siltstone to 6 cm) and are subrounded to subangular. Metasedimentary clasts (gryllite, schist) are similar in shape, but generally smaller in size. Granite and quartz clasts are rounded to well rounded, the latter reaching 30 cm in diameter. Sandstone clasts are dominantly grey-green weathering, fine- to medium-grained, micaceous and feldspathic. In some instances they exhibit postdepositional an echelon tension fractures. Clast counts also indicate that sedimentary clasts are more dominant in this map area compared to more abundant metasedimentary clasts further west. Metasedimentary clasts are dominantly dark grey weathering pelite commonly with an earlier tectonic fabric. The matrix is grey-brown weathering, medium- to very coarse-grained, feldspathic, lithic sandstone. Conglomerates further upsection are finer grained and better organized, displaying imbrication and trough cross-stratification. In exposures near Salmon River Lake, these conglomerates contain a significant amount of plant debris with fragments up to 30 cm long. Sandstone occurs as trough-shaped lenses up to approximately 2 m thick, characteristically 30-80 cm thick, with a minimum lateral extent of 15 m. The lenses are grey-green weathering, medium- to very coarse-grained, feldspathic and micaceous. Occasional interstratified pebbly or granitoides horizons occur. Scours, trough cross-stratification (sets to 30 cm), ripple cross-stratification and normal grading all occur in the sandstone. The number of sandstone lenses increases upsection.

The deposition of coarse conglomerate occurs along the southern flank of the Basin suggesting a strong tectonic influence on sedimentation where subsidence along the Basin margin occurs along northerly dipping tectonic normal faults. In contrast, the character of the sediments does not vary with proximity to the northern margin (Chedabucto Fault) suggesting that the Fault does not constitute the original Basin margin, and that an unknown portion of the Basin and its Meguma basement have been tectonically removed and may be found north of the Fault.

The St. Marys Basin is an example of basin development and evolution adjacent to an intracratonic fault zone associated with oblique convergence during orogenesis. Its evolution provides constraints on the potential relationship between the termination of the mid-Paleozoic Acadian Orogeny, subsequent basin development and the ongoing interactions between the Avalon and Meguma terranes, and between Laurentia and Gondwana during the assembly of Pangaea. More generally, because the relationship between tectonic development and motion along intracratonic strike-slip faults in continental zones is difficult to interpret, the sedimentology and structural geology in basins developed along these fault zones may preserve a less ambiguous record of the main tectonic events.

The Late Paleozoic evolution of the St. Marys Basin, mainland Nova Scotia, preserves evidence of protracted dextral shear along an intracratonic fault zone during collisional orogenesis and the assembly of Pangaea. The St. Marys Basin formed within the eastern extension of the Meguma Fault Zone (MFZ) along the boundary between the Avalon and Meguma terranes and contains latest Devonian-Tournaisian continental clastic rocks that are 3000-4000 m in thickness.

The origin and evolution of the Basin are attributed to either discrete or progressive dextral strike-slip tectonics along the MFZ between the Late Devonian and Late Carboniferous. Evidence for the Late Devonian origin of the Basin is recorded along its southern flank by the fabrics of the deformed ca. 370 Ma granites, the overall sedimentary facies distribution and some syndepositional features within the clastic rocks. The most intense deformation within the Basin is concentrated in a narrow east-northeastward-trending zone, in which predominantly fine grained clastic rocks are deformed into pendular folds and related reverse faults. The orientation of this zone, relative to the MFZ, is consistent with dextral shear. At least some of this deformation occurred after the deposition of the overlying Viséan Windsor Group. The style of deformation along the present northern margin of the Basin (the Chedabucto Fault) is also consistent with regional dextral shear.


## Disclaimer

The information on this map may have come from a variety of government and non-government sources. The Nova Scotia Department of Natural Resources does not assume any liability for errors that may occur.

## Recommended Citation

Murphy, J.B. and Rice, R.J. 2005. Geological map for part of NTS 11F/05, Guysborough area, Nova Scotia; Nova Scotia Department of Natural Resources, Mineral Resources Branch, Open File Map ME 2005-113, scale 1:50 000.

## Regional Key Map



## Map Notes

Universal Transverse Mercator Projection (UTM), Zone 20, Central Meridian 63°00' West, North American Datum (NAD) 1927.  
Base and digital data derived from the Nova Scotia Topographic Database (NSTDB). The NSTDB is available from Service Nova Scotia and Municipal Relations (SNMRL), Land Information Services Division (LIS), Nova Scotia Geomatics Centre (NSGC), Annapolis, Nova Scotia.

## Acknowledgments

We acknowledge the support of the Geological Survey of Canada, Natural Sciences and Engineering Research Council, Caledonia Mining Corporation and the Nova Scotia Department of Natural Resources. We are grateful to David Dove, Samantha Evans, Lorne Jenx, Fraser Keppie, Jon Malone, Sarah Jane Mosher, Cathal Meagher, Alex Nazok, Jacquie Stevens, Tim Stokes and Tim Webster for assistance in the field.

Map production funded by Natural Resources Canada and Nova Scotia Department of Natural Resources under the Targeted Geoscience Initiative (Phase 2) Project: Geological Mapping and Resource Evaluation in Central Nova Scotia, 2004-2005.

Cartography and reproduction by Nova Scotia Department of Natural Resources, Geoscience Information Services Section, 2004-2005.

## LEGEND

### CARBONIFEROUS

#### WINDSOR GROUP

EWG undivided

### LATE DEVONIAN - EARLY CARBONIFEROUS

#### HORTON GROUP

LD-EChet WEST RIVER ST. MARYS FORMATION

redish-brown to grey-brown weathering, pebble to boulder conglomerates interstratified with grey-brown weathering sandstone. Abundant clasts derived from the Meguma Terrane.

LD-EChcb CROSS BROOK FORMATION

grey-green weathering sandstone, interbedded grey-green weathering siltstone, shale, clay, very coarse-grained sandstone, micaceous, rounded or polyhedral granitoides and conglomerate interstratified with red-brown to grey-brown weathering shale, siltstone, sandstone, carbonate and organic debris.

LD-EChgr LD-ECh LD-EChswn

GRAHAM HILLS', LOCHIEL, BARRENS HILLS' and LITTLE STEWACKE RIVER FORMATIONS (LD-EChgr) undivided  
red and maroon weathering, fine grained (fine to felspathic) siltstone and siltstone with thick intervals of grey weathering, interstratified, coarse grained sandstone, pebbly sandstone and granule to pebble conglomerates. Clasts include quartz, mica, metamorphosed siltstone (0.5-1 cm), and flow-banded mudstone.

LD-Chet LD-Chgr LD-EChswn  
BARRENS HILLS AND LITTLE STEWACKE RIVER FORMATIONS (LD-EChgr) undivided  
red and maroon weathering, fine grained (fine to felspathic) siltstone and siltstone with thick intervals of grey weathering, interstratified, coarse grained sandstone, pebbly sandstone and granule to pebble conglomerates. Clasts include quartz, mica, metamorphosed siltstone (0.5-1 cm), and flow-banded mudstone.

LITTLE STEWACKE RIVER FORMATION (LD-EChswn) interstratified, finely bedded, blocky mudstone, shale, silt to dark grey and black shale and silt, and light to dark grey sandstone containing comminuted plant debris.

-bathym units are not shown in this map area

## Geological Symbols

- Outcrop
- Bedding, tops known (inclined, overturn, vertical, horizontal)
- Bedding, tops unknown (inclined, vertical, horizontal)
- Intersection lineation, 1st generation
- Cleavage or foliation, 1st generation (inclined, vertical)
- Minor fold axis
- Anticline and syncline (approximate)
- Minor fault
- Major fault (defined)
- Geological boundary (defined, approximate, assumed)
- Interfingering contact, approximate (units of approximately similar age, units of differing age)

## Legend

- Contour
- Index Contour
- Depression Contour
- Index Depression Contour
- Coastline
- Lakes, Single-line Rivers, Streams
- 100 Series Highway
- Trans-Canada Highway
- Trunk Highway
- Collector Highway
- Hard Surface Road
- Road Under Construction
- Loose Surface/Resource Access Road
- Vehicle Track
- Trail/Footpath
- Railway
- Railway Inactive
- County Boundary