
BARITE, CELESTITE AND FLUORITE

IN NOVA SCOTIA

BY 7.H. FOWLER

Information Circular 15



BARITE, CELESTITE AND FLUORITE

IN NOVA SCOTIA

BY 7.H. FOWLER

Information Circular 15 Second Edition

Nova Scotia Department of Natural Resources Halifax, Nova Scotia 1991



Barite, Celestite and Fluorite

Introduction

Barite, celestite and fluorite are internationally traded commodities. Demand is concentrated in the industrialized countries of Europe, North America and Japan. Supplies are available in most regions of the world.

Approximately 85 per cent of the world production of barite is used as a weighting agent in oil and gas well drilling. The remaining 15 per cent is consumed for the manufacture of barium compounds, as a filler and extender in the manufacture of plastics, rubber, paint, adhesives and sealants, in glass making, ceramics and in pharmaceuticals.

Celestite, the principal source of strontium, is consumed primarily by the chemical industry where it is converted into strontium compounds, mainly strontium carbonate and strontium nitrate. Strontium carbonate is used in colour television picture tubes to shield against the emission of X-rays and to provide a clearer picture. Strontium carbonate is also used in the production of ceramic ferrite magnets, pigments, in the manufacture of glass, in the chemical and electronics industries, in zinc refining and in the manufacture of aluminum castings. The principal use for strontium nitrate is in manufacturing pyrotechnics.

Fluorite, the most important source of fluorine, is used in the manufacture of hydrofluoric acid and fluorine chemicals; as a fluxing agent in steel making; in the electrolytic reduction of alumina to aluminum; in refining of uranium ores; and in the glass and ceramics industries.

Mining History

Barite

Barite mining in Nova Scotia has continued virtually uninterrupted since 1865. The earliest reported mining was at the Eureka Mine at Five Islands. Colchester County in 1865. In 1868 harite mining started at Stewiacke, Colchester County, followed by mining at Lake Ainslie and Cape Rouge, Inverness County in 1894, to produce barite for paint manufacturing. By 1900 Nova Scotia had seven producing barite mines, mostly to supply high-purity product to paint manufacturing plants within the Province. Several of these mines were operated by Henderson and Potts, a paint company hased in Halifax. In 1912, Barytes Ltd. built the Province's first harite grinding mill at Scotsville, Inverness County, to process ore from nearby deposits. Production statistics from 1904 to 1920 show Nova Scotia deposits furnished most of the entire output of Canadian barite. Barite mining continued steadily until 1925, then waned until the discovery of the Walton, Hants County deposit in 1940. In 1951, Maritime Barvtes Co. Ltd. developed a barite deposit at Upper Brookfield, Colchester County and erected a mill to produce high-purity filler grade barite. Inconsistent product grade forced this operation to close in 1952.

The economic possibilities of an occurrence of barite at Walton were not appreciated until 1940, when Springer Gold Mines Limited completed an extensive diamond drilling program which outlined Canada's largest barite mine, the Walton Mine. In 1941 a company was incorporated under the name Canadian

Industrial Minerals Limited to develop the deposit, first as an open-pit mine, and later as an open-pit and underground mining operation. In 1955 Magnet Cove Barium Corporation, a subsidiary of Dresser Industries Inc., Dallas, Texas, purchased the property from Canadian Industrial Minerals Limited, Diamond drilling in 1956-57 discovered a lead-zinccopper-silver ore body below the barite deposit and sulphide production began in 1961. In 1967 Magnet Cove Barium Corporation became Dresser Minerals which operated the mine until it closed in 1978. The Walton Mine produced barite for 38 years. Most of the barite was shipped to United States and Caribbean destinations where it was used as a weighting agent in drilling fluids with smaller amounts shipped to the Middle East for the same purpose. Lesser amounts of chemical and filler grade barite were shipped to Central Canadian markets. During the operating life of the Walton Mine, 4.3 Mt of barite were produced from the open-pit and underground. In 1976 the Mine was placed on a phase-out schedule and on February 15, 1978 production at Walton ceased, leaving an estimated 900 000 t of barite underground.

After the closure of the Walton Mine, barite mining came to an end in the Province until the development in 1980 of the Upper Brookfield barite-siderite deposit and the construction of a processing plant at Debert, Colchester County, by Nystone Chemicals Ltd. to produce pharmaceutical-grade barium sulphate. This was followed by the opening of an open-pit barite mine at Lake Uist, Richmond County in 1983 by Novex Mining and Exploration Company

Limited to produce drill mud grade barite. Two other barite properties were developed to produce drill mud grade barite. Magcohar Dresser Canada mined barite from a small high grade deposit at Bass River, Colchester County in 1984-1985. In 1985 Scotsville Mineral Resources developed the MacInnis barite deposit at Scotsville, Inverness County. Only Nystone Chemicals Ltd. in Upper Brookfield is producing barite in Nova Scotia today.

Celestite

Significant deposits of celestite were discovered by Lura Corporation at Enon, Cape Breton County in 1962 and these low grade deposits were developed by Kaiser Celestite Mining in 1970. A concentrator to upgrade the ore was constructed on site and a chemical plant was built at Point Edward 60 km northeast of the mine. The first shipment of celestite concentrate from the Enon mill to the chemical plant was in 1971. At the Point Edward plant the concentrate was converted to SrCO₃, Sr(NO₃)₂, Na₂SO₄ and Sr(OH)₂. Operations continued until 1976 when production ended because of lost markets to higher grade deposits in Mexico.

Barite Production

Nystone Chemicals Ltd. is the only harite producer in Nova Scotia. Nystone has a barite-siderite deposit in Early Carboniferous sedimentary rocks at Upper Brookfield, Colchester County. This open-pit mine supplies ore to the Nystone processing plant at Debert, Colchester County. The mill circuit at the plant consists of gravity separation,

magnetic separation, acid leaching and ultra fine grinding in a paddle mill to produce USP pharmaceutical grade barium sulphate. This product is shipped to E-Z-EM Inc. in Westbury, New York, U.S.A. where it is prepared and packaged into barium kits which are sold to hospitals and medical clinics.

General Geology of Nova Scotia Barite, Fluorite and Celestite

Nova Scotia has over 70 identified occurrences and deposits of barite, celestite and fluorite. Seventy per cent of these occurrences are hosted by sedimentary rocks, mostly of Early Carboniferous age. Mineralization occurs as fissure fillings, pore space fillings or solution cavity fillings and as replacement deposits. Seventeen per cent of the occurrences are found in igneous rocks. The intrusive host rocks are Devonian age and the extrusive rocks are either Early or Late Carboniferous age. The mineralization is fissure filling type. The remaining 13 per cent of the occurrences are the fissure filling type hosted by metamorphic rocks ranging in age from Precambrian to Silurian (Felderhof, 1978).

Barite-Fluorite Development Potential

Two properties having very good development potential in Nova Scotia are the Scotsville barite deposits at Scotsville, Inverness County and the Conwest barite-fluorite deposits at East Lake Ainslie, Inverness County (Fig. 1). At

Scotsville, drill mud grade barite was produced in 1985 by Scotsville Mineral Resources at the MacInnis open-pit Mine.




Figure 1. Location of major barite, celestite and fluorite deposits.

The Conwest barite-fluorite property has large undeveloped reserves of barite and fluorite. The property is owned by Conwest Exploration Company Limited of Toronto, Ontario.

Scotsville Barite Deposits

The Peter Campbell Mine and the MacInnis Mine (*Photo 1*) occur near Scotsville, Inverness County at the northeastern corner of Lake Ainslie. Access is by Highway 395 which leads



Photo 1. MacInnis Mine, Scotsville, Inverness County, 1985.

from Whycocomagh, Inverness County situated on Trans-Canada Highway 105. These deposits are commonly referred to as the Scotsville barite deposits. Both occur in what is known as the Scotsville inlier, a northward-trending belt composed of Precambrian, Devonian and Early Carboniferous rocks, forming a highland area along the northeastern side of Lake Ainslie.

In 1984 Scotsville Mineral Resources began mining at the MacInnis Mine. Site development work included clear cutting, stripping, road construction, water reservoir development, construction of concrete foundation for ore processing equipment and the establishment of waste disposal sites. The barite orebody occurs at the base of the Lower Carboniferous Horton Group sedimentary rocks, which unconformably overlie George River Group metasedimentary rocks and intrusive rocks of Precambrian age. Most of the barite occurs within the basal Horton breccia which dips from 35° to 60° east. Ore reserves to a depth of 45 m are estimated at 95 000 t having an average grade of 60 per cent BaSO₄. Additional reserves are drill inferred below 45 m. Minor amounts of fluorite and trace amounts of pyrite, chalcopyrite and malachite are the only mineral contaminants and these do not impede the jigging of the barite to mud grade specifications. During the early 1900s, 10 000 t of barite from this deposit were used for paint manufacturing. Approximately 25 000 t of barite are immediately available at the MacInnis deposit requiring a minimum of stripping.

The Peter Campbell Mine and several other barite veins at Scotsville are good exploration targets. There is no barite mining in this area at the present time.

Conwest Barite-Fluorite Deposits

Conwest Exploration Company Limited owns an undeveloped barite-fluorite property at East Lake Ainslie, Inverness County. Access to the property is by Highway 395 which leads from the Trans-Canada Highway 105 at Whycocomagh, Inverness County.

The property has nine barite-fluorite veins with reserves estimated at 4.1 Mt averaging 33.4 per cent barite and 17.4 per cent fluorite. These deposits have been developed along fault cavities and subsidiary tension fractures and shears mostly in rhyolitic tuff of the Early Carboniferous Fisset Brook Formation. One exception is the western ore extension of one of the veins which lies along the fault contact of the tuff and Horton Group sedimentary rocks. Two of these deposits, the Campbell-MacMillan deposit and the Upper Johnson deposit, contain the major tonnage of baritefluorite. The Campbell-MacMillan vein strikes east-west and has a 65° to 70° southerly dip. Reserves have been calculated at 2.6 Mt averaging 27.8 percent barite and 19 per cent fluorite. The Upper Johnson vein (Photo 2) strikes S85°W and dips 85° north to vertical. Reserves are calculated at 960 000 t of 45 per cent barite and 17 per cent fluorite.



Photo 2. Upper Johnson barite-fluorite vein, Conwest property, Inverness County.

Metallurgical studies and pilot plot testing show that mud grade barite and metallurgical grade and acid grade fluorspar can be produced from these deposits. A program is currently under way to determine the suitability of the barite for the manufacture of specialty barite compounds. At present there is no mining activity at any of these deposits.

Barite Occurrences

Nova Scotia has numerous other barite occurrences; many have been small producers in the past. The hest occurrences for prospecting are those found in Windsor Group rocks of Early Carboniferous age found on mainland Nova Scotia and Cape Breton Island. The Pine Brook barite property near the Community of Lake Uist, Richmond County did produce mud grade barite in 1983. This barite occurs as an irregular replacement of limestone meinber in Windsor Group sedimentary rocks. Exploration programs are necessary to establish barite reserves at all of these occurrences.

Fluorite Occurrences

Fluorite, other than the Conwest baritefluorite deposits at Lake Ainslie, occurs at several locations in Devonian age granitic rocks and Lower Carboniferous sedimentary rocks. Most occurrences are found in granitic rocks in the New Ross area of Lunenburg County. All occurrences have been prospected and are considered as having no mining potential.

Celestite Development Potential

The most significant celestite deposits in Nova Scotia are those occurring in the Loch Lomond Sedimentary Basin near the Village of Enon, Cape Breton County. These deposits are known as the Enon, Amac and MacRae deposits (Fig. 1). The Enon and MacRae deposits were mined by Kaiser Celestite Mining from 1970 to 1976. Kaiser mined 272 000 t of ore from these deposits and ran it through their 450 t/day flotation concentrator at Enon. The concentrator ungraded the ore to over 90 per cent celestite prior to shipping to their own chemical plant at Point Edward, Cape Breton County, where it was converted to SrCO₃, Sr(NO₃)₂, Na₂SO₄ and Sr(OH)₂.

MacRae Celestite Deposit

The MacRae Celestite deposit (Fig. 2) has excellent development potential. This deposit is 50 km south of Sydney and 80 km north of the deep water ocean port at the Strait of Canso.




Figure 2. Location of MacRae celestite deposit, Enon, Cape Breton County.

Kaiser Celestite Mining drilled 90 diamond-drill holes in the MacRae deposit and intersected four celestite zones, indicated stratigraphically from bottom to top as zones, 1, 2, 3 and 4 with zones 2 and 3 considered minable. The ore zones are celestite replacement in silty limestone and silty limestone conglomerates. These tabular ore zones dip from 17° to 20° to the northwest and are separated by conglomerate and limestone. Ore zones 2 and 3 contain in excess of 500 000 t of minable celestite with an average grade of over 50 per cent. A summary of the MacRae celestite ore zone is shown in Table 1. It is significant to note that the flotation mill constructed by Kaiser is still standing and could be used to upgrade the MacRae ore (see cover). At the present time Timminco Limited of Toronto holds a special licence on the MacRae deposit and is examining its development potential.

Celestite Occurrences

Along with the celestite deposits at Enon, Cape Breton County, Nova Scotia has five known celestite occurrences. All of these occurrences are hosted by Early Carboniferous age, Windsor Group rocks. Three occurrences are on Cape Breton Island and two are on mainland Nova Scotia. All of these occurrences have been prospected. The Beckwith, Cumberland County occurrence has received the most attention because of its high grade ore. The most recent work at Beckwith was the bulk sampling and metallurgical testing by Timminco Limited in 1986.

Table 1. Summary of celestite ore zones, MacRae deposit*

Thickness Range (m)	Average (m)	Host Rock	Mineralization
2.4–2.7	2.5	Celestite replacing limestone	Celestite, massive and disseminated, fine- to coarse-grained, light grey to grey brown.
0.6-9.1	3.7	Celestite replacing silty limestone	Celestite, massive very dense, brittle, light grey to brown, medium- to coarse-grained.
0.6–7.0	3.1	Celestite replacing silty limestone conglomerate	Celestite, medium- to coarse-grained, vuggy with vugs lined with celestite crystals.
0.3-1.5	0.5	Celestite replacing silty limestone conglomerate	Celestite, medium grained, light hrown.
	Range (m) 2.4–2.7 0.6–9.1 0.6–7.0	Range (m) 2.4–2.7 2.5 0.6–9.1 3.7 0.6–7.0 3.1	Range (m) 2.4–2.7 2.5 Celestite replacing limestone 0.6–9.1 3.7 Celestite replacing silty limestone 0.6–7.0 3.1 Celestite replacing silty limestone 0.3–1.5 O.5 Celestite replacing silty limestone Celestite replacing silty limestone Celestite replacing silty limestone

Mineral Rights

Barite, celestite and fluorite are minerals in Nova Scotia which are reserved to the Crown. A Crown mineral is defined as a natural solid, inorganic or fossilized organic substance and a substance prescribed to be a mineral. A landowner or anyone else may acquire the mineral rights by making application for a mineral exploration licence. The regulations and procedures for acquiring and maintaining a licence are set forth in the Mineral Resources Act, Statutes of Nova Scotia. This Act is administered by the Minister of Natural Resources, Province of Nova Scotia.

Selected Bibliography

- Felderhof, G.W., 1978: Barite, celestite and fluorite in Nova Scotia; Nova Scotia Department of Mines, Bulletin No. 4.
- Forgeron, S., 1977: The Kaiser Celestite Mining operation and mineral potential of the Loch Lomond Basin, Cape Breton, Nova Scotia; Nova Scotia Department of Mines and Energy, Open File Report 328.
- Messervey, J.P., 1930: Barytes in Nova Scotia; Nova Scotia Department of Public Works and Mines, Pamphlet No. 4.
- Zurowski, M., 1972: Barite-fluorite deposits of Lake Ainslie-an appraisal from an economic viewpoint; Canadian Institute of Mining and Mctallurgy Transactions, v. LXXV, p. 318-321.



Department of Natural Resources

Honourable C.W. MacNeil, M.D. Minister

Halifax, Nova Scotia, 1991

Nova Scotia Department of Natural Resources, Information Circular No. 15, Second Edition

Canada–Nova Scotia Cooperation Agreement on Mineral Development

For additional information contact:

Nova Scotia Department of Natural Resources 1701 Hollis Street P.O. Box 698 Halifax, Nova Scotia B3J 2T9

Telephone: 902-424-4161

Fax: 902-424-7735

