

DATA REPORT 7656: Glenholme, NS

Prepared 27 April 2023
by C. Robicheau, Conservation Data
Analyst

CONTENTS OF REPORT

1.0 Preface

- 1.1 Data List
- 1.2 Restrictions
- 1.3 Additional Information
- Map 1: Buffered Study Area

2.0 Rare and Endangered Species

- 2.1 Flora
- 2.2 Fauna
- Map 2: Flora and Fauna

3.0 Special Areas

- 3.1 Managed Areas
- 3.2 Significant Areas
- Map 3: Special Areas

4.0 Rare Species Lists

- 4.1 Fauna
- 4.2 Flora
- 4.3 Location Sensitive Species
- 4.4 Source Bibliography

5.0 Rare Species within 100 km

- 5.1 Source Bibliography



Map 1. A 100 km buffer around the study area

1.0 PREFACE

The Atlantic Canada Conservation Data Centre (AC CDC; www.accdc.com) is part of a network of NatureServe data centres and heritage programs serving 50 states in the U.S.A, 10 provinces and 1 territory in Canada, plus several Central and South American countries. The NatureServe network is more than 30 years old and shares a common conservation data methodology. The AC CDC was founded in 1997, and maintains data for the jurisdictions of New Brunswick, Nova Scotia, Prince Edward Island, and Newfoundland and Labrador. Although a non-governmental agency, the AC CDC is supported by 6 federal agencies and 4 provincial governments, as well as through outside grants and data processing fees.

Upon request and for a fee, the AC CDC queries its database and produces customized reports of the rare and endangered flora and fauna known to occur in or near a specified study area. As a supplement to that data, the AC CDC includes locations of managed areas with some level of protection, and known sites of ecological interest or sensitivity.

1.1 DATA LIST

Included datasets:

| <u>Filename</u> | <u>Contents</u> |
|-----------------------------|---|
| GlenholmeNS_7656ob.xls | Rare or legally-protected Flora and Fauna in your study area |
| GlenholmeNS_7656ob100km.xls | A list of Rare and legally protected Flora and Fauna within 100 km of your study area |
| GlenholmeNS_7656msa.xls | Managed and Biologically Significant Areas in your study area |
| GlenholmeNS_7656ff_py.xls | Rare Freshwater Fish in your study area (DFO database) |

1.2 RESTRICTIONS

The AC CDC makes a strong effort to verify the accuracy of all the data that it manages, but it shall not be held responsible for any inaccuracies in data that it provides. By accepting AC CDC data, recipients assent to the following limits of use:

- a) Data is restricted to use by trained personnel who are sensitive to landowner interests and to potential threats to rare and/or endangered flora and fauna posed by the information provided.
- b) Data is restricted to use by the specified Data User; any third party requiring data must make its own data request.
- c) The AC CDC requires Data Users to cease using and delete data 12 months after receipt, and to make a new request for updated data if necessary at that time.
- d) AC CDC data responses are restricted to the data in our Data System at the time of the data request.
- e) Each record has an estimate of locational uncertainty, which must be referenced in order to understand the record's relevance to a particular location. Please see attached Data Dictionary for details.
- f) AC CDC data responses are not to be construed as exhaustive inventories of taxa in an area.
- g) The absence of a taxon cannot be inferred by its absence in an AC CDC data response.

1.3 ADDITIONAL INFORMATION

The accompanying Data Dictionary provides metadata for the data provided.

Please direct any additional questions about AC CDC data to the following individuals:

Plants, Lichens, Ranking Methods, All other Inquiries

Sean Blaney
Senior Scientist / Executive Director
(506) 364-2658
sean.blaney@accdc.ca

Animals (Fauna)

John Klymko
Zoologist
(506) 364-2660
john.klymko@accdc.ca

Data Management, GIS

James Churchill
Conservation Data Analyst / Field Biologist
(902) 679-6146
james.churchill@accdc.ca

Billing

Jean Breau
Financial Manager / Executive Assistant
(506) 364-2657
jean.breau@accdc.ca

Questions on the biology of Federal Species at Risk can be directed to AC CDC: (506) 364-2658, with questions on Species at Risk regulations to: Samara Eaton, Canadian Wildlife Service (NB and PE): (506) 364-5060 or Julie McKnight, Canadian Wildlife Service (NS): (902) 426-4196.

For provincial information about rare taxa and protected areas, or information about game animals, deer yards, old growth forests, archeological sites, fish habitat etc., in New Brunswick, please contact Hubert Askanas, Energy and Resource Development: (506) 453-5873.

For provincial information about rare taxa and protected areas, or information about game animals, deer yards, old growth forests, archeological sites, fish habitat etc., in Nova Scotia, please contact Donna Hurlburt, NS DLF: (902) 679-6886. To determine if location-sensitive species (section 4.3) occur near your study site please contact a NS DLF Regional Biologist:

Western: Emma Vost
(902) 670-8187
Emma.Vost@novascotia.ca

Western: Sarah Spencer
(902) 541-0081
Sarah.Spencer@novascotia.ca

Central: Shavonne Meyer
(902) 893-0816
Shavonne.Meyer@novascotia.ca

Central: Kimberly George
(902) 890-1046
Kimberly.George@novascotia.ca

Eastern: Harrison Moore
(902) 497-4119
Harrison.Moore@novascotia.ca

Eastern: Maureen Cameron-MacMillan
(902) 295-2554
Maureen.Cameron-MacMillan@novascotia.ca

Eastern: Elizabeth Walsh
(902) 563-3370
Elizabeth.Walsh@novascotia.ca

For provincial information about rare taxa and protected areas, or information about game animals, fish habitat etc., in Prince Edward Island, please contact Garry Gregory, PEI Dept. of Communities, Land and Environment: (902) 569-7595.

2.0 RARE AND ENDANGERED SPECIES


2.1 FLORA

The study area contains 4 records of 3 vascular and 1 record of 1 nonvascular flora (Map 2 and attached: *ob.xls), excluding 'location-sensitive' species.

2.2 FAUNA

The study area contains 191 records of 40 vertebrate and 11 records of 5 invertebrate fauna (Map 2 and attached data files - see 1.1 Data List), excluding 'location-sensitive' species. Please see section 4.3 to determine if 'location-sensitive' species occur near your study site.

Map 2: Known observations of rare and/or protected flora and fauna within the study area.



- RESOLUTION**
- 4.7 within 50s of kilometers
 - 4.0 within 10s of kilometers
 - 3.7 within 5s of kilometers
 - △ 3.0 within kilometers
 - △ 2.7 within 500s of meters
 - ◇ 2.0 within 100s of meters
 - ◇ 1.7 within 10s of meters

- HIGHER TAXON**
- vertebrate fauna
 - invertebrate fauna
 - vascular flora
 - nonvascular flora

3.0 SPECIAL AREAS


3.1 MANAGED AREAS

The GIS scan identified 1 managed area in the vicinity of the study area (Map 3 and attached file: *ma*.xls).

3.2 SIGNIFICANT AREAS

The GIS scan identified 3 biologically significant sites in the vicinity of the study area (Map 3 and attached file: *sa*.xls).

Map 3: Boundaries and/or locations of known Managed and Significant Areas within the study area.



4.0 RARE SPECIES LISTS

Rare and/or endangered taxa (excluding “location-sensitive” species, section 4.3) within the study area listed in order of concern, beginning with legally listed taxa, with the number of observations per taxon and the distance in kilometers from study area centroid to the closest observation (\pm the precision, in km, of the record). [P] = vascular plant, [N] = nonvascular plant, [A] = vertebrate animal, [I] = invertebrate animal, [C] = community. Note: records are from attached files *ob.xls/*ob.shp only.

4.1 FLORA

| | Scientific Name | Common Name | COSEWIC | SARA | Prov Legal Prot | Prov Rarity Rank | # recs | Distance (km) |
|---|---------------------------|---------------------|-----------------|-----------------|-----------------|------------------|--------|---------------|
| N | <i>Pectenia plumbea</i> | Blue Felt Lichen | Special Concern | Special Concern | Vulnerable | S3 | 1 | 2.3 \pm 5.0 |
| P | <i>Lobelia spicata</i> | Pale-Spiked Lobelia | | | | S1 | 1 | 2.6 \pm 7.0 |
| P | <i>Juncus subcaudatus</i> | Woods-Rush | | | | S3S4 | 2 | 0.5 \pm 5.0 |
| P | <i>Huperzia appressa</i> | Mountain Firmoss | | | | S3S4 | 1 | 2.6 \pm 7.0 |

4.2 FAUNA

| | Scientific Name | Common Name | COSEWIC | SARA | Prov Legal Prot | Prov Rarity Rank | # recs | Distance (km) |
|---|--------------------------------|---|-----------------|-----------------|-----------------|------------------|--------|---------------|
| A | <i>Salmo salar pop. 1</i> | Atlantic Salmon - Inner Bay of Fundy population | Endangered | Endangered | | S1 | 2 | 3.3 \pm 0.0 |
| A | <i>Riparia riparia</i> | Bank Swallow | Threatened | Threatened | Endangered | S2B | 15 | 0.8 \pm 1.0 |
| A | <i>Limosa haemastica</i> | Hudsonian Godwit | Threatened | | | S2S3M | 1 | 2.1 \pm 0.0 |
| A | <i>Anguilla rostrata</i> | American Eel | Threatened | | | S3N | 2 | 3.3 \pm 0.0 |
| A | <i>Phalaropus lobatus</i> | Red-necked Phalarope | Special Concern | Special Concern | | S2S3M | 1 | 2.1 \pm 0.0 |
| A | <i>Hirundo rustica</i> | Barn Swallow | Special Concern | Threatened | Endangered | S3B | 8 | 1.6 \pm 0.0 |
| A | <i>Cardellina canadensis</i> | Canada Warbler | Special Concern | Threatened | Endangered | S3B | 3 | 1.6 \pm 0.0 |
| A | <i>Chordeiles minor</i> | Common Nighthawk | Special Concern | Threatened | Threatened | S3B | 1 | 2.6 \pm 7.0 |
| A | <i>Contopus cooperi</i> | Olive-sided Flycatcher | Special Concern | Threatened | Threatened | S3B | 4 | 0.7 \pm 0.0 |
| A | <i>Dolichonyx oryzivorus</i> | Bobolink | Special Concern | Threatened | Vulnerable | S3B | 3 | 2.6 \pm 7.0 |
| A | <i>Contopus virens</i> | Eastern Wood-Pewee | Special Concern | Special Concern | Vulnerable | S3S4B | 1 | 2.6 \pm 7.0 |
| A | <i>Sialia sialis</i> | Eastern Bluebird | Not At Risk | | | S3B | 2 | 1.2 \pm 0.0 |
| A | <i>Ammospiza nelsoni</i> | Nelson's Sparrow | Not At Risk | | | S3S4B | 5 | 1.8 \pm 0.0 |
| A | <i>Calidris canutus rufa</i> | Red Knot rufa subspecies | E,SC | Endangered | Endangered | S2M | 3 | 1.1 \pm 0.0 |
| A | <i>Passerina cyanea</i> | Indigo Bunting | | | | S1?B,SUM | 3 | 0.3 \pm 0.0 |
| A | <i>Charadrius semipalmatus</i> | Semipalmated Plover | | | | S1B,S4M | 27 | 1.1 \pm 0.0 |
| A | <i>Calidris minutilla</i> | Least Sandpiper | | | | S1B,S4M | 10 | 1.1 \pm 0.0 |
| A | <i>Molothrus ater</i> | Brown-headed Cowbird | | | | S2B | 2 | 2.6 \pm 7.0 |
| A | <i>Spatula clypeata</i> | Northern Shoveler | | | | S2B,SUM | 2 | 0.9 \pm 0.0 |
| A | <i>Calidris alba</i> | Sanderling | | | | S2N,S3M | 21 | 1.1 \pm 0.0 |
| A | <i>Rallus limicola</i> | Virginia Rail | | | | S2S3B | 5 | 0.9 \pm 0.0 |
| A | <i>Spinus pinus</i> | Pine Siskin | | | | S3 | 1 | 2.6 \pm 7.0 |
| A | <i>Salvelinus fontinalis</i> | Brook Trout | | | | S3 | 6 | 3.3 \pm 0.0 |
| A | <i>Spatula discors</i> | Blue-winged Teal | | | | S3B | 1 | 2.6 \pm 7.0 |
| A | <i>Charadrius vociferus</i> | Killdeer | | | | S3B | 6 | 1.1 \pm 0.0 |
| A | <i>Tringa semipalmata</i> | Willet | | | | S3B | 2 | 2.6 \pm 7.0 |
| A | <i>Tyrannus tyrannus</i> | Eastern Kingbird | | | | S3B | 3 | 2.6 \pm 7.0 |
| A | <i>Pheucticus ludovicianus</i> | Rose-breasted Grosbeak | | | | S3B | 1 | 2.6 \pm 7.0 |
| A | <i>Falco sparverius</i> | American Kestrel | | | | S3B,S4S5M | 1 | 2.6 \pm 7.0 |
| A | <i>Gallinago delicata</i> | Wilson's Snipe | | | | S3B,S5M | 2 | 2.6 \pm 7.0 |
| A | <i>Setophaga tigrina</i> | Cape May Warbler | | | | S3B,SUM | 1 | 2.6 \pm 7.0 |
| A | <i>Pluvialis squatarola</i> | Black-bellied Plover | | | | S3M | 5 | 1.1 \pm 0.0 |
| A | <i>Arenaria interpres</i> | Ruddy Turnstone | | | | S3M | 1 | 1.1 \pm 0.0 |
| A | <i>Calidris pusilla</i> | Semipalmated Sandpiper | | | | S3M | 28 | 1.1 \pm 0.0 |
| A | <i>Limnodromus griseus</i> | Short-billed Dowitcher | | | | S3M | 3 | 2.1 \pm 0.0 |
| A | <i>Picoides arcticus</i> | Black-backed Woodpecker | | | | S3S4 | 1 | 2.6 \pm 7.0 |
| A | <i>Botaurus lentiginosus</i> | American Bittern | | | | S3S4B,S4S5M | 2 | 2.6 \pm 7.0 |

| | Scientific Name | Common Name | COSEWIC | SARA | Prov Legal Prot | Prov Rarity Rank | # recs | Distance (km) |
|---|----------------------------------|-------------------------|---------|------|-----------------|------------------|--------|---------------|
| A | <i>Actitis macularia</i> | Spotted Sandpiper | | | | S3S4B,S5M | 3 | 2.6 ± 7.0 |
| A | <i>Leiothlypis peregrina</i> | Tennessee Warbler | | | | S3S4B,S5M | 2 | 2.6 ± 7.0 |
| A | <i>Eremophila alpestris</i> | Horned Lark | | | | SHB,S4S5N,S5M | 1 | 2.6 ± 7.0 |
| I | <i>Coenagrion resolutum</i> | Taiga Bluet | | | | S2 | 1 | 0.9 ± 0.0 |
| I | <i>Aglais milberti</i> | Milbert's Tortoiseshell | | | | S2S3 | 6 | 1.6 ± 2.0 |
| I | <i>Polygonia interrogationis</i> | Question Mark | | | | S3B | 1 | 4.5 ± 2.0 |
| I | <i>Cecropia pylades</i> | Northern Cloudywing | | | | S3S4 | 1 | 4.5 ± 2.0 |
| I | <i>Polygonia faunus</i> | Green Comma | | | | S3S4 | 2 | 1.6 ± 2.0 |

4.3 LOCATION SENSITIVE SPECIES

The Department of Natural Resources in each Maritimes province considers a number of species “location sensitive”. Concern about exploitation of location-sensitive species precludes inclusion of precise coordinates in this report. Those intersecting your study area are indicated below with “YES”.

Nova Scotia

| Scientific Name | Common Name | SARA | Prov Legal Prot | Known within the Study Site? |
|---|---|---------------------------|---------------------------|------------------------------|
| <i>Fraxinus nigra</i> | Black Ash | | Threatened | No |
| <i>Emydoidea blandingii</i> | Blanding's Turtle - Nova Scotia pop. | Endangered | Endangered | No |
| <i>Glyptemys insculpta</i> | Wood Turtle | Threatened | Threatened | No |
| <i>Falco peregrinus pop. 1</i> | Peregrine Falcon - anatum/tundrius pop. | | Vulnerable | No |
| <i>Bat hibernaculum</i> or bat species occurrence | | [Endangered] ¹ | [Endangered] ¹ | No |

¹ *Myotis lucifugus* (Little Brown Myotis), *Myotis septentrionalis* (Long-eared Myotis), and *Perimyotis subflavus* (Tri-colored Bat or Eastern Pipistrelle) are all Endangered under the Federal Species at Risk Act and the NS Endangered Species Act.

4.4 SOURCE BIBLIOGRAPHY

The recipient of these data shall acknowledge the AC CDC and the data sources listed below in any documents, reports, publications or presentations, in which this dataset makes a significant contribution.

| # recs | CITATION |
|--------|---|
| 101 | Morrison, Guy. 2011. Maritime Shorebird Survey (MSS) database. Canadian Wildlife Service, Ottawa, 15939 surveys. 86171 recs. |
| 47 | Lepage, D. 2014. Maritime Breeding Bird Atlas Database. Bird Studies Canada, Sackville NB, 407,838 recs. |
| 24 | Erskine, A.J. 1992. Maritime Breeding Bird Atlas Database. NS Museum & Nimbus Publ., Halifax, 82,125 recs. |
| 10 | iNaturalist. 2020. iNaturalist Data Export 2020. iNaturalist.org and iNaturalist.ca, Web site: 128728 recs. |
| 8 | eBird. 2020. eBird Basic Dataset. Version: EBD_relNov-2019. Ithaca, New York. Nov 2019, Cape Breton Bras d'Or Lakes Watershed subset. Cornell Lab of Ornithology. |
| 4 | Webster, R.P. Atlantic Forestry Centre Insect Collection, Maritimes butterfly records. Natural Resources Canada. 2014. |
| 3 | Amiro, P.G. 1998. Atlantic Salmon Inner Bay of Fundy SFA 22 & part of 23. DFO Sci. SSR D3-12. |
| 3 | Klymko, J. Dataset of butterfly records at the New Brunswick Museum not yet accessioned by the museum. Atlantic Canada Conservation Data Centre. 2016. |
| 2 | Benjamin, L.K. (compiler). 2012. Significant Habitat & Species Database. Nova Scotia Dept Natural Resources, 4965 recs. |
| 2 | Zinck, M. & Roland, A.E. 1998. Roland's Flora of Nova Scotia. Nova Scotia Museum, 3rd ed., rev. M. Zinck; 2 Vol., 1297 pp. |
| 1 | Bird Studies Canada. 2020. Important Bird and Biodiversity Areas in Canada database (Retrieved: 28 July, 2020 from https://www.ibacanada.com/explore.jsp?lang=EN). IBA Program. |
| 1 | Brunelle, P.-M. (compiler). 2009. ADIP/MDDS Odonata Database: data to 2006 inclusive. Atlantic Dragonfly Inventory Program (ADIP), 24200 recs. |
| 1 | Clayden, S. Digitization of Wolfgang Maass Nova Scotia forest lichen collections, 1964-2004. New Brunswick Museum. 2018. |
| 1 | Ferguson, D.C. 1954. The Lepidoptera of Nova Scotia. Part I, macrolepidoptera. Proceedings of the Nova Scotian Institute of Science, 23(3), 161-375. |
| 1 | Hill, N.M. 1994. Status report on the Long's bulrush <i>Scirpus longii</i> in Canada. Committee on the Status of Endangered Wildlife in Canada, 7 recs. |
| 1 | Layberry, R.A. & Hall, P.W., LaFontaine, J.D. 1998. The Butterflies of Canada. University of Toronto Press. 280 pp+plates. |
| 1 | Newell, R.E. 2005. E.C. Smith Digital Herbarium. E.C. Smith Herbarium, Irving Biodiversity Collection, Acadia University, Web site: http://luxor.acadiau.ca/library/Herbarium/project/ . 582 recs. |
| 1 | Nova Scotia Dept Natural Resources, Forestry Branch. 2007. Restricted & Limited Use Land Database (RLUL). , http://www.gov.ns.ca/natr/FORESTRY/rlul/downloadrlul.htm . |
| 1 | Ogden, J. NS DNR Butterfly Collection Dataset. Nova Scotia Department of Natural Resources. 2014. |
| 1 | Porter, C.J.M. 2014. Field work data 2007-2014. Nova Scotia Nature Trust, 96 recs. |

5.0 RARE SPECIES WITHIN 100 KM

A 100 km buffer around the study area contains 52673 records of 145 vertebrate and 1822 records of 72 invertebrate fauna; 8549 records of 287 vascular and 2900 records of 189 nonvascular flora (attached: *ob100km.xls).

Taxa within 100 km of the study site that are rare and/or endangered in the province in which the study site occurs (including “location-sensitive” species). All ranks correspond to the province in which the study site falls, even for out-of-province records. Taxa are listed in order of concern, beginning with legally listed taxa, with the number of observations per taxon and the distance in kilometers from study area centroid to the closest observation (\pm the precision, in km, of the record).

| Taxonomic Group | Scientific Name | Common Name | COSEWIC | SARA | Prov Legal Prot | Prov Rarity Rank | # recs | Distance (km) | Prov |
|-----------------|---|--|-----------------|-----------------|-----------------|------------------|--------|-----------------|------|
| A | <i>Coregonus huntsmani</i> | Atlantic Whitefish | Endangered | Endangered | Endangered | S1 | 1 | 77.5 \pm 1.0 | NS |
| A | <i>Myotis lucifugus</i> | Little Brown Myotis | Endangered | Endangered | Endangered | S1 | 183 | 12.6 \pm 0.0 | NS |
| A | <i>Myotis septentrionalis</i> | Northern Myotis | Endangered | Endangered | Endangered | S1 | 23 | 12.6 \pm 0.0 | NS |
| A | <i>Perimyotis subflavus</i> | Tricolored Bat | Endangered | Endangered | Endangered | S1 | 13 | 14.2 \pm 1.0 | NS |
| A | <i>Salmo salar pop. 1</i> | Atlantic Salmon - Inner Bay of Fundy population | Endangered | Endangered | | S1 | 50 | 3.3 \pm 0.0 | NS |
| A | <i>Salmo salar pop. 6</i> | Atlantic Salmon - Nova Scotia Southern Upland population | Endangered | | | S1 | 25 | 58.6 \pm 0.0 | NS |
| A | <i>Charadrius melodus melodus</i> | Piping Plover melodus subspecies | Endangered | Endangered | Endangered | S1B | 870 | 28.5 \pm 0.0 | NS |
| A | <i>Sterna dougallii</i> | Roseate Tern | Endangered | Endangered | Endangered | S1B | 25 | 83.9 \pm 0.0 | NS |
| A | <i>Dermochelys coriacea pop. 2</i> | Leatherback Sea Turtle - Atlantic population | Endangered | Endangered | | S1S2N | 3 | 81.3 \pm 1.0 | NB |
| A | <i>Morone saxatilis pop. 2</i> | Striped Bass - Bay of Fundy population | Endangered | | | S2S3B,S2S3N | 4 | 44.5 \pm 0.0 | NS |
| A | <i>Catharus bicknelli</i> | Bicknell's Thrush | Threatened | Threatened | Endangered | S1B | 1 | 62.1 \pm 7.0 | NS |
| A | <i>Asio flammeus</i> | Short-eared Owl | Threatened | Special Concern | | S1B | 52 | 48.9 \pm 7.0 | NS |
| A | <i>Glyptemys insculpta</i> | Wood Turtle | Threatened | Threatened | Threatened | S2 | 1046 | 8.7 \pm 5.0 | NS |
| A | <i>Riparia riparia</i> | Bank Swallow | Threatened | Threatened | Endangered | S2B | 2611 | 0.8 \pm 1.0 | NS |
| A | <i>Chaetura pelagica</i> | Chimney Swift | Threatened | Threatened | Endangered | S2S3B,S1M | 1312 | 8.2 \pm 0.0 | NS |
| A | <i>Limosa haemastica</i> | Hudsonian Godwit | Threatened | | | S2S3M | 203 | 2.1 \pm 0.0 | NS |
| A | <i>Acipenser oxyrinchus</i> | Atlantic Sturgeon | Threatened | | | S2S3N | 7 | 31.9 \pm 0.0 | NS |
| A | <i>Hydrobates leucorhous</i> | Leach's Storm-Petrel | Threatened | | | S3B | 6 | 84.4 \pm 0.0 | NS |
| A | <i>Tringa flavipes</i> | Lesser Yellowlegs | Threatened | | | S3M | 1613 | 7.9 \pm 0.0 | NS |
| A | <i>Anguilla rostrata</i> | American Eel | Threatened | | | S3N | 54 | 3.3 \pm 0.0 | NS |
| A | <i>Sturnella magna</i> | Eastern Meadowlark | Threatened | Threatened | | SHB | 4 | 60.8 \pm 7.0 | NS |
| A | <i>Ixobrychus exilis</i> | Least Bittern | Threatened | Threatened | | SUB | 12 | 69.6 \pm 0.0 | NS |
| A | <i>Hylocichla mustelina</i> | Wood Thrush | Threatened | Threatened | | SUB | 55 | 18.3 \pm 7.0 | NS |
| A | <i>Salmo salar pop. 12</i> | Atlantic Salmon - Gaspé - Southern Gulf of St. Lawrence population | Special Concern | | | S1 | 31 | 31.1 \pm 50.0 | NS |
| A | <i>Antrostomus vociferus</i> | Eastern Whip-Poor-Will | Special Concern | Threatened | Threatened | S1?B | 15 | 44.0 \pm 7.0 | NS |
| A | <i>Passerculus sandwichensis princeps</i> | Ipswich Sparrow | Special Concern | Special Concern | | S1B | 3 | 84.1 \pm 0.0 | NS |
| A | <i>Bucephala islandica</i> | Barrow's Goldeneye | Special Concern | Special Concern | | S1N,SUM | 11 | 54.0 \pm 1.0 | NS |
| A | <i>Euphagus carolinus</i> | Rusty Blackbird | Special Concern | Special Concern | Endangered | S2B | 259 | 19.7 \pm 0.0 | NS |
| A | <i>Phalaropus lobatus</i> | Red-necked Phalarope | Special Concern | Special Concern | | S2S3M | 15 | 2.1 \pm 0.0 | NS |
| A | <i>Histrionicus histrionicus pop. 1</i> | Harlequin Duck - Eastern population | Special Concern | Special Concern | Endangered | S2S3N,SUM | 15 | 74.0 \pm 0.0 | NS |
| A | <i>Chelydra serpentina</i> | Snapping Turtle | Special Concern | Special Concern | Vulnerable | S3 | 135 | 12.6 \pm 0.0 | NS |
| A | <i>Hirundo rustica</i> | Barn Swallow | Special Concern | Threatened | Endangered | S3B | 1834 | 1.6 \pm 0.0 | NS |
| A | <i>Cardellina canadensis</i> | Canada Warbler | Special Concern | Threatened | Endangered | S3B | 1246 | 1.6 \pm 0.0 | NS |
| A | <i>Chordeiles minor</i> | Common Nighthawk | Special Concern | Threatened | Threatened | S3B | 473 | 2.6 \pm 7.0 | NS |
| A | <i>Contopus cooperi</i> | Olive-sided Flycatcher | Special Concern | Threatened | Threatened | S3B | 1096 | 0.7 \pm 0.0 | NS |
| A | <i>Dolichonyx oryzivorus</i> | Bobolink | Special Concern | Threatened | Vulnerable | S3B | 2011 | 2.6 \pm 7.0 | NS |

| Taxonomic Group | Scientific Name | Common Name | COSEWIC | SARA | Prov Legal Prot | Prov Rarity Rank | # recs | Distance (km) | Prov |
|-----------------|-----------------------------------|------------------------------------|-----------------|-----------------|-----------------|------------------|--------|---------------|------|
| A | <i>Coccothraustes vespertinus</i> | Evening Grosbeak | Special Concern | Special Concern | Vulnerable | S3B,S3N,S3M | 801 | 6.1 ± 0.0 | NS |
| A | <i>Podiceps auritus</i> | Horned Grebe | Special Concern | Special Concern | | S3N,SUM | 19 | 83.8 ± 0.0 | NB |
| A | <i>Contopus virens</i> | Eastern Wood-Pewee | Special Concern | Special Concern | Vulnerable | S3S4B | 1205 | 2.6 ± 7.0 | NS |
| A | <i>Phocoena phocoena</i> | Harbour Porpoise | Special Concern | | | S4 | 3 | 82.4 ± 0.0 | NS |
| A | <i>Chrysemys picta picta</i> | Eastern Painted Turtle | Special Concern | Special Concern | | S4 | 220 | 7.9 ± 0.0 | NS |
| A | <i>Accipiter cooperii</i> | Cooper's Hawk | Not At Risk | | | S1?B,SUN,SUM | 6 | 61.0 ± 7.0 | NS |
| A | <i>Fulica americana</i> | American Coot | Not At Risk | | | S1B | 51 | 17.9 ± 0.0 | NS |
| A | <i>Chlidonias niger</i> | Black Tern | Not At Risk | | | S1B | 178 | 70.0 ± 0.0 | NS |
| A | <i>Falco peregrinus pop. 1</i> | Peregrine Falcon - anatum/tundrius | Not At Risk | Special Concern | Vulnerable | S1B,SUM | 263 | 41.8 ± 0.0 | NS |
| A | <i>Sorex dispar</i> | Long-tailed Shrew | Not At Risk | | | S2 | 2 | 16.5 ± 0.0 | NS |
| A | <i>Aegolius funereus</i> | Boreal Owl | Not At Risk | | | S2?B,SUM | 9 | 11.6 ± 7.0 | NS |
| A | <i>Lynx canadensis</i> | Canada Lynx | Not At Risk | | Endangered | S2S3 | 2 | 94.8 ± 1.0 | NB |
| A | <i>Globicephala melas</i> | Long-finned Pilot Whale | Not At Risk | | | S2S3 | 2 | 61.6 ± 100.0 | NS |
| A | <i>Hemidactylium scutatum</i> | Four-toed Salamander | Not At Risk | | | S3 | 28 | 36.2 ± 0.0 | NS |
| A | <i>Megaptera novaeangliae</i> | Humpback Whale | Not At Risk | | | S3 | 1 | 60.6 ± 0.0 | NS |
| A | <i>Sterna hirundo</i> | Common Tern | Not At Risk | | | S3B | 389 | 39.7 ± 0.0 | NS |
| A | <i>Sialia sialis</i> | Eastern Bluebird | Not At Risk | | | S3B | 144 | 1.2 ± 0.0 | NS |
| A | <i>Buteo lagopus</i> | Rough-legged Hawk | Not At Risk | | | S3N | 16 | 72.3 ± 0.0 | NS |
| A | <i>Accipiter gentilis</i> | Northern Goshawk | Not At Risk | | | S3S4 | 166 | 14.3 ± 0.0 | NS |
| A | <i>Glaucomys volans</i> | Southern Flying Squirrel | Not At Risk | | | S3S4 | 6 | 69.4 ± 10.0 | NS |
| A | <i>Lagenorhynchus acutus</i> | Atlantic White-sided Dolphin | Not At Risk | | | S3S4 | 7 | 51.2 ± 0.0 | NS |
| A | <i>Ammodramus nelsoni</i> | Nelson's Sparrow | Not At Risk | | | S3S4B | 870 | 1.8 ± 0.0 | NS |
| A | <i>Calidris canutus rufa</i> | Red Knot rufa subspecies | E,SC | Endangered | Endangered | S2M | 579 | 1.1 ± 0.0 | NS |
| A | <i>Morone saxatilis</i> | Striped Bass | E,SC | | | S2S3B,S2S3N | 8 | 18.4 ± 0.0 | NS |
| A | <i>Alces alces americana</i> | Moose | | | Endangered | S1 | 207 | 9.8 ± 3.0 | NS |
| A | <i>Picoides dorsalis</i> | American Three-toed Woodpecker | | | | S1? | 2 | 56.2 ± 0.0 | NS |
| A | <i>Uria aalge</i> | Common Murre | | | | S1?B | 1 | 84.5 ± 0.0 | NS |
| A | <i>Passerina cyanea</i> | Indigo Bunting | | | | S1?B,SUM | 36 | 0.3 ± 0.0 | NS |
| A | <i>Nycticorax nycticorax</i> | Black-crowned Night-heron | | | | S1B | 2 | 84.5 ± 1.0 | NB |
| A | <i>Oxyura jamaicensis</i> | Ruddy Duck | | | | S1B | 55 | 70.7 ± 7.0 | NS |
| A | <i>Gallinula galeata</i> | Common Gallinule | | | | S1B | 44 | 40.6 ± 7.0 | NS |
| A | <i>Myiarchus crinitus</i> | Great Crested Flycatcher | | | | S1B | 36 | 23.3 ± 7.0 | NS |
| A | <i>Cistothorus palustris</i> | Marsh Wren | | | | S1B | 56 | 69.8 ± 0.0 | NS |
| A | <i>Mimus polyglottos</i> | Northern Mockingbird | | | | S1B | 96 | 31.8 ± 7.0 | NS |
| A | <i>Toxostoma rufum</i> | Brown Thrasher | | | | S1B | 21 | 13.7 ± 0.0 | NS |
| A | <i>Charadrius semipalmatus</i> | Semipalmated Plover | | | | S1B,S4M | 2073 | 1.1 ± 0.0 | NS |
| A | <i>Calidris minutilla</i> | Least Sandpiper | | | | S1B,S4M | 1432 | 1.1 ± 0.0 | NS |
| A | <i>Anas acuta</i> | Northern Pintail | | | | S1B,SUM | 114 | 8.4 ± 7.0 | NS |
| A | <i>Vireo gilvus</i> | Warbling Vireo | | | | S1B,SUM | 40 | 33.5 ± 7.0 | NS |
| A | <i>Vespertilionidae sp.</i> | bat species | | | | S1S2 | 185 | 5.5 ± 0.0 | NS |
| A | <i>Pooecetes gramineus</i> | Vesper Sparrow | | | | S1S2B,SUM | 80 | 18.3 ± 7.0 | NS |
| A | <i>Vireo philadelphicus</i> | Philadelphia Vireo | | | | S2?B,SUM | 105 | 20.0 ± 7.0 | NS |
| A | <i>Fratercula arctica</i> | Atlantic Puffin | | | | S2B | 1 | 84.9 ± 0.0 | NB |
| A | <i>Empidonax traillii</i> | Willow Flycatcher | | | | S2B | 62 | 12.6 ± 0.0 | NS |
| A | <i>Molothrus ater</i> | Brown-headed Cowbird | | | | S2B | 238 | 2.6 ± 7.0 | NS |
| A | <i>Spatula clypeata</i> | Northern Shoveler | | | | S2B,SUM | 248 | 0.9 ± 0.0 | NS |
| A | <i>Mareca strepera</i> | Gadwall | | | | S2B,SUM | 305 | 21.7 ± 7.0 | NS |
| A | <i>Piranga olivacea</i> | Scarlet Tanager | | | | S2B,SUM | 39 | 12.6 ± 0.0 | NS |
| A | <i>Calidris alba</i> | Sanderling | | | | S2N,S3M | 1528 | 1.1 ± 0.0 | NS |
| A | <i>Asio otus</i> | Long-eared Owl | | | | S2S3 | 28 | 29.3 ± 7.0 | NS |
| A | <i>Rallus limicola</i> | Virginia Rail | | | | S2S3B | 262 | 0.9 ± 0.0 | NS |
| A | <i>Rissa tridactyla</i> | Black-legged Kittiwake | | | | S2S3B | 2 | 27.7 ± 0.0 | NS |
| A | <i>Petrochelidon pyrrhonota</i> | Cliff Swallow | | | | S2S3B | 536 | 11.9 ± 7.0 | NS |
| A | <i>Phalacrocorax carbo</i> | Great Cormorant | | | | S2S3B,S2S3 | 115 | 71.8 ± 3.0 | NS |

| Taxonomic Group | Scientific Name | Common Name | COSEWIC | SARA | Prov Legal Prot | Prov Rarity Rank | # recs | Distance (km) | Prov |
|-----------------|-------------------------------------|-------------------------|---------|------|-----------------|----------------------|--------|---------------|------|
| A | <i>Cathartes aura</i> | Turkey Vulture | | | | N S2S3B,S4S5 M | 127 | 44.8 ± 0.0 | NS |
| A | <i>Setophaga pinus</i> | Pine Warbler | | | | S2S3B,S4S5 M | 55 | 8.4 ± 7.0 | NS |
| A | <i>Bucephala clangula</i> | Common Goldeneye | | | | S2S3B,S5N,S 5M | 147 | 10.9 ± 5.0 | NS |
| A | <i>Icterus galbula</i> | Baltimore Oriole | | | | S2S3B,SUM | 97 | 7.6 ± 0.0 | NS |
| A | <i>Pluvialis dominica</i> | American Golden-Plover | | | | S2S3M | 197 | 7.9 ± 0.0 | NS |
| A | <i>Numerius phaeopus hudsonicus</i> | Whimbrel | | | | S2S3M | 239 | 18.3 ± 0.0 | NS |
| A | <i>Phalaropus fulicarius</i> | Red Phalarope | | | | S2S3M | 4 | 82.7 ± 0.0 | NS |
| A | <i>Perisoreus canadensis</i> | Canada Jay | | | | S3 | 724 | 8.4 ± 7.0 | NS |
| A | <i>Poecile hudsonicus</i> | Boreal Chickadee | | | | S3 | 770 | 5.9 ± 0.0 | NS |
| A | <i>Spinus pinus</i> | Pine Siskin | | | | S3 | 625 | 2.6 ± 7.0 | NS |
| A | <i>Salvelinus fontinalis</i> | Brook Trout | | | | S3 | 68 | 3.3 ± 0.0 | NS |
| A | <i>Salvelinus namaycush</i> | Lake Trout | | | | S3 | 2 | 55.9 ± 0.0 | NS |
| A | <i>Sorex maritimensis</i> | Maritime Shrew | | | | S3 | 105 | 82.3 ± 1.0 | NB |
| A | <i>Synaptomys cooperi</i> | Southern Bog Lemming | | | | S3 | 1 | 67.9 ± 0.0 | NS |
| A | <i>Pekania pennanti</i> | Fisher | | | | S3 | 6 | 40.4 ± 0.0 | NS |
| A | <i>Calcarius lapponicus</i> | Lapland Longspur | | | | S3?N,SUM | 34 | 82.7 ± 2.0 | NB |
| A | <i>Spatula discors</i> | Blue-winged Teal | | | | S3B | 470 | 2.6 ± 7.0 | NS |
| A | <i>Charadrius vociferus</i> | Killdeer | | | | S3B | 987 | 1.1 ± 0.0 | NS |
| A | <i>Tringa semipalmata</i> | Willet | | | | S3B | 1404 | 2.6 ± 7.0 | NS |
| A | <i>Sterna paradisaea</i> | Arctic Tern | | | | S3B | 27 | 82.2 ± 0.0 | NB |
| A | <i>Coccyzus erythrophthalmus</i> | Black-billed Cuckoo | | | | S3B | 128 | 8.4 ± 7.0 | NS |
| A | <i>Tyrannus tyrannus</i> | Eastern Kingbird | | | | S3B | 472 | 2.6 ± 7.0 | NS |
| A | <i>Pheucticus ludovicianus</i> | Rose-breasted Grosbeak | | | | S3B | 683 | 2.6 ± 7.0 | NS |
| A | <i>Alosa pseudoharengus</i> | Alewife | | | | S3B | 30 | 16.9 ± 0.0 | NS |
| A | <i>Somateria mollissima</i> | Common Eider | | | | S3B,S3M,S3N | 327 | 41.8 ± 7.0 | NS |
| A | <i>Tringa melanoleuca</i> | Greater Yellowlegs | | | | S3B,S4M | 2481 | 7.9 ± 0.0 | NS |
| A | <i>Falco sparverius</i> | American Kestrel | | | | S3B,S4S5M | 569 | 2.6 ± 7.0 | NS |
| A | <i>Gallinago delicata</i> | Wilson's Snipe | | | | S3B,S5M | 1402 | 2.6 ± 7.0 | NS |
| A | <i>Setophaga striata</i> | Blackpoll Warbler | | | | S3B,S5M | 132 | 11.6 ± 7.0 | NS |
| A | <i>Cardellina pusilla</i> | Wilson's Warbler | | | | S3B,S5M | 126 | 8.4 ± 7.0 | NS |
| A | <i>Pinicola enucleator</i> | Pine Grosbeak | | | | S3B,S5N,S5M | 145 | 11.6 ± 7.0 | NS |
| A | <i>Setophaga tigrina</i> | Cape May Warbler | | | | S3B,SUM | 330 | 2.6 ± 7.0 | NS |
| A | <i>Branta bernicla</i> | Brant | | | | S3M | 22 | 59.0 ± 0.0 | NS |
| A | <i>Pluvialis squatarola</i> | Black-bellied Plover | | | | S3M | 2002 | 1.1 ± 0.0 | NS |
| A | <i>Arenaria interpres</i> | Ruddy Turnstone | | | | S3M | 702 | 1.1 ± 0.0 | NS |
| A | <i>Calidris pusilla</i> | Semipalmated Sandpiper | | | | S3M | 2445 | 1.1 ± 0.0 | NS |
| A | <i>Calidris melanotos</i> | Pectoral Sandpiper | | | | S3M | 437 | 7.9 ± 0.0 | NS |
| A | <i>Limnodromus griseus</i> | Short-billed Dowitcher | | | | S3M | 1517 | 2.1 ± 0.0 | NS |
| A | <i>Chroicocephalus ridibundus</i> | Black-headed Gull | | | | S3N | 11 | 77.2 ± 7.0 | NS |
| A | <i>Picoides arcticus</i> | Black-backed Woodpecker | | | | S3S4 | 198 | 2.6 ± 7.0 | NS |
| A | <i>Loxia curvirostra</i> | Red Crossbill | | | | S3S4 | 218 | 8.4 ± 7.0 | NS |
| A | <i>Sorex palustris</i> | American Water Shrew | | | | S3S4 | 2 | 82.3 ± 1.0 | NB |
| A | <i>Botaurus lentiginosus</i> | American Bittern | | | | S3S4B,S4S5 M | 1016 | 2.6 ± 7.0 | NS |
| A | <i>Setophaga castanea</i> | Bay-breasted Warbler | | | | S3S4B,S4S5 M | 738 | 5.9 ± 0.0 | NS |
| A | <i>Actitis macularius</i> | Spotted Sandpiper | | | | S3S4B,S5M | 1006 | 2.6 ± 7.0 | NS |
| A | <i>Leiophyips peregrina</i> | Tennessee Warbler | | | | S3S4B,S5M | 670 | 2.6 ± 7.0 | NS |
| A | <i>Passerella iliaca</i> | Fox Sparrow | | | | S3S4B,S5M | 71 | 11.9 ± 7.0 | NS |
| A | <i>Mergus serrator</i> | Red-breasted Merganser | | | | S3S4B,S5M,S 5N | 206 | 20.1 ± 0.0 | NS |
| A | <i>Calidris maritima</i> | Purple Sandpiper | | | | S3S4N | 90 | 50.6 ± 15.0 | NS |
| A | <i>Lanius borealis</i> | Northern Shrike | | | | S3S4N | 27 | 57.5 ± 0.0 | NS |

| Taxonomic Group | Scientific Name | Common Name | COSEWIC | SARA | Prov Legal Prot | Prov Rarity Rank | # recs | Distance (km) | Prov |
|-----------------|---|----------------------------------|-----------------|-----------------|-----------------|------------------|--------|---------------|------|
| A | <i>Morus bassanus</i> | Northern Gannet | | | | SHB | 74 | 56.5 ± 4.0 | NS |
| A | <i>Aythya americana</i> | Redhead | | | | SHB | 7 | 70.7 ± 7.0 | NS |
| A | <i>Leucophaeus atricilla</i> | Laughing Gull | | | | SHB | 14 | 80.2 ± 0.0 | NB |
| A | <i>Progne subis</i> | Purple Martin | | | | SHB | 24 | 36.6 ± 7.0 | NS |
| A | <i>Eremophila alpestris</i> | Horned Lark | | | | SHB,S4S5N,S5M | 21 | 2.6 ± 7.0 | NS |
| I | <i>Bombus bohemicus</i> | Ashton Cuckoo Bumble Bee | Endangered | Endangered | Endangered | S1 | 38 | 8.9 ± 5.0 | NS |
| I | <i>Danaus plexippus</i> | Monarch | Endangered | Special Concern | Endangered | S2?B,S3M | 507 | 13.0 ± 0.0 | NS |
| I | <i>Danaus plexippus plexippus</i> | Monarch | Endangered | Special Concern | | S2?B,S3M | 1 | 60.0 ± 0.0 | NS |
| I | <i>Barnea truncata</i> | Atlantic Mud-piddock | Threatened | Threatened | | S1 | 1 | 18.1 ± 1.0 | NS |
| I | <i>Bombus suckleyi</i> | Suckley's Cuckoo Bumble Bee | Threatened | | | SH | 1 | 21.5 ± 5.0 | NS |
| I | <i>Alasmidonta varicosa</i> | Brook Floater | Special Concern | Special Concern | Threatened | S3 | 11 | 29.6 ± 0.0 | NS |
| I | <i>Bombus terricola</i> | Yellow-banded Bumble Bee | Special Concern | Special Concern | Vulnerable | S3 | 126 | 8.9 ± 5.0 | NS |
| I | <i>Coccinella transversoguttata richardsoni</i> | Transverse Lady Beetle | Special Concern | | Endangered | SH | 9 | 15.0 ± 2.0 | NS |
| I | <i>Gomphurus ventricosus</i> | Skillet Clubtail | Special Concern | Endangered | | SH | 2 | 51.3 ± 0.0 | NS |
| I | <i>Cicindela formosa</i> | Big Sand Tiger Beetle | | | | S1 | 1 | 81.2 ± 1.0 | NS |
| I | <i>Erora laeta</i> | Early Hairstreak | | | | S1 | 2 | 83.9 ± 1.0 | NS |
| I | <i>Pachydiplax longipennis</i> | Blue Dasher | | | | S1 | 4 | 76.7 ± 0.0 | NS |
| I | <i>Atlanticoncha ochracea</i> | Tidewater Mucket | | | | S1 | 29 | 56.5 ± 0.0 | NS |
| I | <i>Polygonia comma</i> | Eastern Comma | | | | S1? | 19 | 71.2 ± 0.0 | NS |
| I | <i>Polygonia satyrus</i> | Satyr Comma | | | | S1? | 9 | 25.7 ± 5.0 | NS |
| I | <i>Euphyes bimacula</i> | Two-spotted Skipper | | | | S1S2 | 3 | 80.5 ± 0.0 | NS |
| I | <i>Boloria chariclea</i> | Arctic Fritillary | | | | S1S2 | 10 | 26.6 ± 2.0 | NS |
| I | <i>Somatochlora brevicincta</i> | Quebec Emerald | | | | S1S2 | 2 | 67.0 ± 1.0 | NS |
| I | <i>Tharsalea dospassosi</i> | Maritime Copper | | | | S2 | 84 | 42.8 ± 0.0 | NS |
| I | <i>Satyrrium acadica</i> | Acadian Hairstreak | | | | S2 | 14 | 8.0 ± 2.0 | NS |
| I | <i>Neurocordulia michaeli</i> | Broad-tailed Shadowdragon | | | | S2 | 9 | 76.8 ± 0.0 | NS |
| I | <i>Coenagrion resolutum</i> | Taiga Bluet | | | | S2 | 11 | 0.9 ± 0.0 | NS |
| I | <i>Margaritifera margaritifera</i> | Eastern Pearlshell | | | | S2 | 130 | 11.3 ± 0.0 | NS |
| I | <i>Pantala hymenaea</i> | Spot-Winged Glider | | | | S2?B | 7 | 67.0 ± 1.0 | NS |
| I | <i>Nymphalis l-album</i> | Compton Tortoiseshell | | | | S2S3 | 17 | 8.7 ± 2.0 | NS |
| I | <i>Aglais milberti</i> | Milbert's Tortoiseshell | | | | S2S3 | 24 | 1.6 ± 2.0 | NS |
| I | <i>Aglais milberti milberti</i> | Milbert's Tortoise Shell | | | | S2S3 | 5 | 47.9 ± 0.0 | NS |
| I | <i>Lanthus vernalis</i> | Southern Pygmy Clubtail | | | | S2S3 | 4 | 92.4 ± 0.0 | NS |
| I | <i>Somatochlora kennedyi</i> | Kennedy's Emerald | | | | S2S3 | 6 | 60.2 ± 1.0 | NS |
| I | <i>Somatochlora williamsoni</i> | Williamson's Emerald | | | | S2S3 | 6 | 77.7 ± 0.0 | NS |
| I | <i>Williamsonia fletcheri</i> | Ebony Boghaunter | | | | S2S3 | 11 | 37.8 ± 0.0 | NS |
| I | <i>Stylurus scudderi</i> | Zebra Clubtail | | | | S2S3 | 6 | 47.2 ± 1.0 | NS |
| I | <i>Alasmidonta undulata</i> | Triangle Floater | | | | S2S3 | 33 | 34.7 ± 0.0 | NS |
| I | <i>Astyleiopus variegatus</i> | Variiegated Long-horned Beetle | | | | S3 | 1 | 75.2 ± 0.0 | NS |
| I | <i>Strophiona nitens</i> | Chestnut Bark Long-horned Beetle | | | | S3 | 2 | 79.9 ± 0.0 | NS |
| I | <i>Hippodamia parenthesis</i> | Parenthesis Lady Beetle | | | | S3 | 2 | 86.1 ± 0.0 | NS |
| I | <i>Naemia seriata</i> | Seaside Lady Beetle | | | | S3 | 21 | 61.5 ± 0.0 | NS |
| I | <i>Chilocorus stigma</i> | Twice-stabbed Lady Beetle | | | | S3 | 2 | 86.5 ± 0.0 | NS |
| I | <i>Myzia pullata</i> | Streaked Lady Beetle | | | | S3 | 1 | 89.4 ± 0.0 | NB |
| I | <i>Monochamus marmorator</i> | Balsam Fir Sawyer | | | | S3 | 1 | 90.5 ± 0.0 | NS |
| I | <i>Trachysida aspera</i> | Rough Flower Longhorn Beetle | | | | S3 | 1 | 76.3 ± 0.0 | NS |
| I | <i>Dicerca tenebrosa</i> | Dark Jewel Beetle | | | | S3 | 1 | 95.6 ± 0.0 | NS |
| I | <i>Astylopsis sexguttata</i> | Six-speckled Long-horned Beetle | | | | S3 | 1 | 65.6 ± 0.0 | NS |
| I | <i>Satyrrium calanus</i> | Banded Hairstreak | | | | S3 | 64 | 8.0 ± 2.0 | NS |
| I | <i>Callophrys lanoraieensis</i> | Bog Elfin | | | | S3 | 12 | 41.3 ± 1.0 | NS |
| I | <i>Strymon melinus</i> | Gray Hairstreak | | | | S3 | 5 | 46.0 ± 2.0 | NS |

| Taxonomic Group | Scientific Name | Common Name | COSEWIC | SARA | Prov Legal Prot | Prov Rarity Rank | # recs | Distance (km) | Prov |
|-----------------|--|---|-----------------|-----------------|-----------------|------------------|--------|---------------|------|
| I | <i>Phanogomphus descriptus</i> | Harpoon Clubtail | | | | S3 | 4 | 32.6 ± 0.0 | NS |
| I | <i>Ophiogomphus aspersus</i> | Brook Snaketail | | | | S3 | 6 | 43.5 ± 0.0 | NS |
| I | <i>Ophiogomphus mainensis</i> | Maine Snaketail | | | | S3 | 15 | 66.5 ± 0.0 | NS |
| I | <i>Ophiogomphus rupinsulensis</i> | Rusty Snaketail | | | | S3 | 28 | 51.2 ± 0.0 | NS |
| I | <i>Epithea princeps</i> | Prince Baskettail | | | | S3 | 17 | 33.1 ± 1.0 | NS |
| I | <i>Somatochlora forcipata</i> | Forcinate Emerald | | | | S3 | 4 | 82.7 ± 1.0 | NS |
| I | <i>Enallagma vernale</i> | Vernal Bluet | | | | S3 | 5 | 16.5 ± 1.0 | NS |
| I | <i>Strophitus undulatus</i> | Creeper | | | | S3 | 6 | 44.8 ± 0.0 | NS |
| I | <i>Polygonia interrogationis</i> | Question Mark | | | | S3B | 219 | 4.5 ± 2.0 | NS |
| I | <i>Cecropiterus pylades</i> | Northern Cloudywing | | | | S3S4 | 18 | 4.5 ± 2.0 | NS |
| I | <i>Amblyscirtes hegon</i> | Pepper and Salt Skipper | | | | S3S4 | 30 | 19.3 ± 0.0 | NS |
| I | <i>Cupido comyntas</i> | Eastern Tailed Blue | | | | S3S4 | 21 | 12.2 ± 0.0 | NS |
| I | <i>Argynnis aphrodite</i> | Aphrodite Fritillary | | | | S3S4 | 43 | 10.6 ± 2.0 | NS |
| I | <i>Polygonia faunus</i> | Green Comma | | | | S3S4 | 20 | 1.6 ± 2.0 | NS |
| I | <i>Oeneis jutta</i> | Jutta Arctic | | | | S3S4 | 34 | 41.1 ± 1.0 | NS |
| I | <i>Aeshna clepsydra</i> | Mottled Darner | | | | S3S4 | 10 | 47.5 ± 0.0 | NS |
| I | <i>Aeshna constricta</i> | Lance-Tipped Darner | | | | S3S4 | 31 | 19.9 ± 1.0 | NS |
| I | <i>Boyeria grafiana</i> | Ocellated Darner | | | | S3S4 | 4 | 31.6 ± 0.0 | NS |
| I | <i>Gomphaeschna furcillata</i> | Harlequin Darner | | | | S3S4 | 7 | 41.2 ± 1.0 | NS |
| I | <i>Somatochlora franklini</i> | Delicate Emerald | | | | S3S4 | 5 | 35.1 ± 1.0 | NS |
| I | <i>Nannothemis bella</i> | Elfin Skimmer | | | | S3S4 | 29 | 41.5 ± 1.0 | NS |
| I | <i>Sympetrum danae</i> | Black Meadowhawk | | | | S3S4 | 1 | 98.5 ± 1.0 | PE |
| I | <i>Amphiagrion saucium</i> | Eastern Red Damsel | | | | S3S4 | 2 | 19.9 ± 1.0 | NS |
| I | <i>Sphaerophoria pyrrhina</i> | Violaceous Globetail | | | | SH | 1 | 22.9 ± 5.0 | NS |
| I | <i>Icaricia saepiolus</i> | Greenish Blue | | | | SH | 4 | 40.5 ± 5.0 | NS |
| I | <i>Polygonia gracilis</i> | Hoary Comma | | | | SH | 2 | 22.6 ± 2.0 | NS |
| N | <i>Erioderma mollissimum</i> | Graceful Felt Lichen | Endangered | Endangered | Endangered | S1 | 13 | 64.7 ± 0.0 | NS |
| N | <i>Erioderma pedicellatum</i> (Atlantic pop.) | Boreal Felt Lichen - Atlantic pop. | Endangered | Endangered | Endangered | S1 | 376 | 64.4 ± 0.0 | NS |
| N | <i>Peltigera hydrothyria</i> | Eastern Waterfan | Threatened | Threatened | Threatened | S1 | 93 | 14.8 ± 0.0 | NS |
| N | <i>Pannaria lurida</i> | Wrinkled Shingle Lichen | Threatened | Threatened | Threatened | S2S3 | 132 | 34.9 ± 0.0 | NS |
| N | <i>Anzia colpododes</i> | Black-foam Lichen | Threatened | Threatened | Threatened | S3 | 38 | 29.4 ± 0.0 | NS |
| N | <i>Fuscopannaria leucosticta</i> | White-rimmed Shingle Lichen | Threatened | | | S3 | 8 | 64.3 ± 0.0 | NS |
| N | <i>Heterodermia squamulosa</i> | Scaly Fringe Lichen | Threatened | | | S3 | 78 | 64.9 ± 0.0 | NS |
| N | <i>Pectenium plumbea</i> | Blue Felt Lichen | Special Concern | Special Concern | Vulnerable | S3 | 184 | 2.3 ± 5.0 | NS |
| N | <i>Sclerophora peronella</i> (Atlantic pop.) | Frosted Glass-whiskers (Atlantic population) | Special Concern | Special Concern | | S3S4 | 26 | 42.0 ± 0.0 | NS |
| N | <i>Pseudevernia cladonia</i> | Ghost Antler Lichen | Not At Risk | | | S2S3 | 9 | 70.6 ± 0.0 | NS |
| N | <i>Fissidens exilis</i> | Pygmy Pocket Moss | Not At Risk | | | S3 | 15 | 25.9 ± 0.0 | NS |
| N | <i>Chaenotheca servitii</i> | Flexuous Golden Stubble | Data Deficient | | | S1 | 1 | 77.7 ± 1.0 | NS |
| N | <i>Erioderma pedicellatum</i> | Boreal Felt Lichen | E,SC | | Endangered | S1 | 1 | 72.4 ± 0.0 | NS |
| N | <i>Aloina brevirostris</i> | Short-Beaked Rigid Screw Moss | | | | S1 | 1 | 61.0 ± 2.0 | NS |
| N | <i>Sematophyllum demissum</i> | a Moss | | | | S1 | 1 | 64.0 ± 2.0 | NS |
| N | <i>Tetradontium brownianum</i> | Little Georgia | | | | S1 | 1 | 64.4 ± 0.0 | NS |
| N | <i>Cyrto-hypnum minutulum</i> | Tiny Cedar Moss | | | | S1 | 1 | 77.9 ± 0.0 | NS |
| N | <i>Blennothallia crispa</i> | Crinkled Jelly Lichen | | | | S1 | 1 | 43.6 ± 0.0 | NS |
| N | <i>Umbilicaria vellea</i> | Grizzled Rocktripe Lichen | | | | S1 | 1 | 80.9 ± 5.0 | NS |
| N | <i>Usnea perplexans</i> | Powdered Beard Lichen | | | | S1 | 1 | 55.2 ± 0.0 | NS |
| N | <i>Scytinium dactylinum</i> | Brown-buttoned Jellyskin Lichen | | | | S1 | 1 | 82.1 ± 0.0 | NS |
| N | <i>Lathagrium cristatum</i> | Fingered Jelly Lichen | | | | S1 | 3 | 58.3 ± 0.0 | NS |
| N | <i>Ephebe perspinulosa</i> | Thread Lichen | | | | S1 | 1 | 82.2 ± 1.0 | NS |
| N | <i>Fuscopannaria praetermissa</i> | Moss Shingles Lichen | | | | S1 | 1 | 63.8 ± 0.0 | NS |
| N | <i>Scytinium schraderi</i> | Wrinkled Jellyskin Lichen | | | | S1 | 1 | 16.8 ± 0.0 | NS |
| N | <i>Lichina confinis</i> | Marine Seaweed Lichen | | | | S1 | 2 | 96.4 ± 0.0 | NS |
| N | <i>Polychidium muscicola</i> | Eyed Mossthorns | | | | S1 | 1 | 58.4 ± 0.0 | NS |

| Taxonomic Group | Scientific Name | Common Name | COSEWIC | SARA | Prov Legal Prot | Prov Rarity Rank | # recs | Distance (km) | Prov |
|-----------------|---------------------------------------|-------------------------------|---------|------|-----------------|------------------|--------|---------------|------|
| N | <i>Sticta limbata</i> | Woollybear Lichen | | | | S1 | 1 | 98.4 ± 3.0 | NS |
| N | <i>Peltigera lepidophora</i> | Powdered Moon Lichen | | | | S1 | 6 | 35.3 ± 4.0 | NS |
| N | <i>Bryoria nitidula</i> | Scaly Pelt Lichen | | | | S1 | 2 | 98.6 ± 0.0 | NS |
| N | <i>Hypogymnia hultenii</i> | Tundra Horsehair Lichen | | | | S1 | 10 | 84.0 ± 0.0 | NS |
| N | <i>Calypogeia neogaea</i> | Powdered Honeycomb Lichen | | | | S1? | 1 | 35.1 ± 0.0 | NS |
| N | <i>Aloina rigida</i> | Common Pouchwort | | | | S1? | 5 | 26.6 ± 0.0 | NS |
| N | <i>Imbricaria muehlenbeckii</i> | Aloe-Like Rigid Screw Moss | | | | S1? | 2 | 81.8 ± 0.0 | NS |
| N | <i>Campylostelium saxicola</i> | Muehlenbeck's Bryum Moss | | | | S1? | 1 | 92.1 ± 0.0 | PE |
| N | <i>Tortula obtusifolia</i> | a Moss | | | | S1? | 3 | 21.2 ± 1.0 | NS |
| N | <i>Didymodon tophaceus</i> | Olive Beard Moss | | | | S1? | 2 | 43.6 ± 4.0 | NS |
| N | <i>Paludella squarrosa</i> | Tufted Fen Moss | | | | S1? | 3 | 53.5 ± 0.0 | NS |
| N | <i>Physcomitrium immersum</i> | a Moss | | | | S1? | 1 | 72.5 ± 0.0 | NS |
| N | <i>Schistostega pennata</i> | Luminous Moss | | | | S1? | 1 | 47.7 ± 0.0 | NS |
| N | <i>Trichodon cylindricus</i> | Cylindric Hairy-teeth Moss | | | | S1? | 1 | 98.1 ± 3.0 | NS |
| N | <i>Syntrichia ruralis</i> | a Moss | | | | S1? | 1 | 69.1 ± 0.0 | NS |
| N | <i>Enchylium limosum</i> | Lime-loving Tarpaper Lichen | | | | S1? | 1 | 43.6 ± 4.0 | NS |
| N | <i>Scytinium intermedium</i> | Forty-five Jellyskin Lichen | | | | S1? | 1 | 43.6 ± 4.0 | NS |
| N | <i>Arrhenopterum heterostichum</i> | One-sided Groove Moss | | | | S1S2 | 2 | 38.1 ± 1.0 | NS |
| N | <i>Brachythecium turgidum</i> | Thick Ragged Moss | | | | S1S2 | 3 | 96.5 ± 0.0 | NS |
| N | <i>Hypnum pratense</i> | Meadow Plait Moss | | | | S1S2 | 1 | 92.8 ± 3.0 | NS |
| N | <i>Mnium thomsonii</i> | Thomson's Leafy Moss | | | | S1S2 | 1 | 62.0 ± 2.0 | NS |
| N | <i>Tortula acaulon</i> | Cuspidate Earth Moss | | | | S1S2 | 1 | 90.4 ± 2.0 | NS |
| N | <i>Plagiothecium latebricola</i> | Alder Silk Moss | | | | S1S2 | 1 | 43.8 ± 3.0 | NS |
| N | <i>Platydictya confervoides</i> | a Moss | | | | S1S2 | 1 | 63.1 ± 0.0 | NS |
| N | <i>Seligeria donniana</i> | Donian Beardless Moss | | | | S1S2 | 1 | 63.9 ± 3.0 | NS |
| N | <i>Sematophyllum marylandicum</i> | a Moss | | | | S1S2 | 2 | 52.7 ± 6.0 | NS |
| N | <i>Timmia megapolitana</i> | Metropolitan Timmia Moss | | | | S1S2 | 3 | 19.7 ± 0.0 | NS |
| N | <i>Tortula mucronifolia</i> | Mucronate Screw Moss | | | | S1S2 | 1 | 62.0 ± 3.0 | NS |
| N | <i>Pseudotaxiphyllum distichaceum</i> | a Moss | | | | S1S2 | 3 | 49.3 ± 0.0 | NS |
| N | <i>Haplocladium microphyllum</i> | Tiny-leaved Haplocladium Moss | | | | S1S2 | 1 | 9.4 ± 5.0 | NS |
| N | <i>Enchylium bachmanianum</i> | Bachman's Jelly Lichen | | | | S1S2 | 1 | 58.8 ± 0.0 | NS |
| N | <i>Placidium squamulosum</i> | Limy Soil Stipplescale Lichen | | | | S1S2 | 1 | 55.0 ± 6.0 | NS |
| N | <i>Peltigera ponojensis</i> | Pale-bellied Pelt Lichen | | | | S1S2 | 1 | 62.5 ± 0.0 | NS |
| N | <i>Pilophorus cereolus</i> | Powdered Matchstick Lichen | | | | S1S2 | 1 | 28.6 ± 3.0 | NS |
| N | <i>Parmotrema reticulatum</i> | Netted Ruffle Lichen | | | | S1S2 | 5 | 68.4 ± 0.0 | NS |
| N | <i>Parmeliella parvula</i> | Poor-man's Shingles Lichen | | | | S1S2 | 8 | 69.1 ± 0.0 | NS |
| N | <i>Heterodermia galactophylla</i> | Branching Fringe Lichen | | | | S1S3 | 2 | 43.3 ± 0.0 | NS |
| N | <i>Xylopsora friesii</i> | a Lichen | | | | S1S3 | 2 | 85.4 ± 0.0 | NS |
| N | <i>Stereocaulon grande</i> | Grand Foam Lichen | | | | S1S3 | 1 | 13.3 ± 0.0 | NS |
| N | <i>Stereocaulon intermedium</i> | Pacific Brain Foam Lichen | | | | S1S3 | 2 | 71.7 ± 2.0 | NS |
| N | <i>Anacamptodon splachnoides</i> | a Moss | | | | S2 | 2 | 43.8 ± 3.0 | NS |
| N | <i>Sphagnum platyphyllum</i> | Flat-leaved Peat Moss | | | | S2 | 2 | 60.9 ± 3.0 | NS |
| N | <i>Sphagnum subnitens</i> | Lustrous Peat Moss | | | | S2 | 1 | 88.0 ± 2.0 | NS |
| N | <i>Usnea flavocardia</i> | Blood-splattered Beard Lichen | | | | S2 | 1 | 85.9 ± 4.0 | NS |
| N | <i>Cystocoleus ebeneus</i> | Rockgossamer Lichen | | | | S2 | 3 | 75.9 ± 0.0 | NS |
| N | <i>Hypotrachyna catawbiensis</i> | Powder-tipped Antler Lichen | | | | S2 | 1 | 94.6 ± 0.0 | NS |
| N | <i>Scytinium imbricatum</i> | Scaly Jellyskin Lichen | | | | S2 | 1 | 30.8 ± 4.0 | NS |
| N | <i>Nephroma arcticum</i> | Arctic Kidney Lichen | | | | S2 | 2 | 80.0 ± 0.0 | NS |
| N | <i>Nephroma resupinatum</i> | a lichen | | | | S2 | 11 | 69.6 ± 1.0 | NS |
| N | <i>Placynthium flabelliforme</i> | Scaly Ink Lichen | | | | S2 | 1 | 57.6 ± 17.0 | NS |

| Taxonomic Group | Scientific Name | Common Name | COSEWIC | SARA | Prov Legal Prot | Prov Rarity Rank | # recs | Distance (km) | Prov |
|-----------------|--|---------------------------------|---------|------|-----------------|------------------|--------|---------------|------|
| N | <i>Riccardia multifida</i> | Delicate Germanderwort | | | | S2? | 1 | 64.5 ± 0.0 | NS |
| N | <i>Anomodon viticulosus</i> | a Moss | | | | S2? | 1 | 30.5 ± 5.0 | NS |
| N | <i>Weissia muhlenbergiana</i> | a Moss | | | | S2? | 5 | 62.0 ± 1.0 | NS |
| N | <i>Atrichum angustatum</i> | Lesser Smoothcap Moss | | | | S2? | 2 | 18.9 ± 5.0 | NS |
| N | <i>Ptychostomum pendulum</i> | Drooping Bryum | | | | S2? | 1 | 61.0 ± 2.0 | NS |
| N | <i>Drepanocladus polygamus</i> | Polygamous Hook Moss | | | | S2? | 3 | 43.6 ± 4.0 | NS |
| N | <i>Pseudocampyllum radicale</i> | Long-stalked Fine Wet Moss | | | | S2? | 1 | 92.8 ± 3.0 | NS |
| N | <i>Dicranum condensatum</i> | Condensed Broom Moss | | | | S2? | 2 | 92.8 ± 3.0 | NS |
| N | <i>Ditrichum rhynchostegium</i> | a Moss | | | | S2? | 2 | 78.4 ± 1.0 | NS |
| N | <i>Kiaeria starkei</i> | Starke's Fork Moss | | | | S2? | 1 | 78.2 ± 10.0 | NS |
| N | <i>Orthotrichum anomalum</i> | Anomalous Bristle Moss | | | | S2? | 1 | 68.1 ± 2.0 | NS |
| N | <i>Philonotis marchica</i> | a Moss | | | | S2? | 2 | 30.3 ± 0.0 | NS |
| N | <i>Physcomitrium collenchymatum</i> | a Moss | | | | S2? | 1 | 98.1 ± 0.0 | NS |
| N | <i>Platydictya jungermannioides</i> | False Willow Moss | | | | S2? | 1 | 41.6 ± 0.0 | NS |
| N | <i>Saelania glaucescens</i> | Blue Dew Moss | | | | S2? | 1 | 35.8 ± 0.0 | NS |
| N | <i>Cyrtomnium hymenophylloides</i> | Short-pointed Lantern Moss | | | | S2? | 2 | 35.8 ± 0.0 | NS |
| N | <i>Platylomella lescurii</i> | a Moss | | | | S2? | 4 | 46.2 ± 0.0 | NS |
| N | <i>Phyllicum demangeonii</i> | Black Rock-wafer Lichen | | | | S2? | 1 | 76.7 ± 0.0 | NS |
| N | <i>Oxyrrhynchium hians</i> | Light Beaked Moss | | | | S2S3 | 4 | 50.6 ± 25.0 | NS |
| N | <i>Platydictya subtilis</i> | Bark Willow Moss | | | | S2S3 | 3 | 43.8 ± 3.0 | NS |
| N | <i>Plagiomnium rostratum</i> | Long-beaked Leafy Moss | | | | S2S3 | 1 | 96.9 ± 2.0 | NS |
| N | <i>Scorpidium revolvens</i> | Limprichtia Moss | | | | S2S3 | 1 | 53.5 ± 0.0 | NS |
| N | <i>Moelleropsis nebulosa</i> | Blue-gray Moss Shingle Lichen | | | | S2S3 | 48 | 24.6 ± 3.0 | NS |
| N | <i>Moelleropsis nebulosa ssp. frullaniae</i> | Blue-gray Moss Shingle Lichen | | | | S2S3 | 2 | 64.5 ± 0.0 | NS |
| N | <i>Ramalina thrausta</i> | Angelhair Ramalina Lichen | | | | S2S3 | 14 | 35.5 ± 0.0 | NS |
| N | <i>Collema leptaleum</i> | Crumpled Bat's Wing Lichen | | | | S2S3 | 64 | 19.7 ± 0.0 | NS |
| N | <i>Usnea ceratina</i> | Warty Beard Lichen | | | | S2S3 | 1 | 38.3 ± 0.0 | NS |
| N | <i>Usnea hirta</i> | Bristly Beard Lichen | | | | S2S3 | 2 | 84.1 ± 0.0 | NS |
| N | <i>Usnea rubicunda</i> | Red Beard Lichen | | | | S2S3 | 3 | 22.7 ± 0.0 | NS |
| N | <i>Ahtiana aurescens</i> | Eastern Candlewax Lichen | | | | S2S3 | 13 | 50.2 ± 0.0 | NS |
| N | <i>Usnocetraria oakesiana</i> | Yellow Band Lichen | | | | S2S3 | 11 | 54.7 ± 0.0 | NS |
| N | <i>Cladonia mateocyatha</i> | Mixed-up Pixie-cup | | | | S2S3 | 3 | 77.2 ± 6.0 | NS |
| N | <i>Cladonia parasitica</i> | Fence-rail Lichen | | | | S2S3 | 4 | 13.3 ± 1.0 | NS |
| N | <i>Chaenotheca gracilentia</i> | a lichen | | | | S2S3 | 1 | 84.4 ± 0.0 | NS |
| N | <i>Scytinium tenuissimum</i> | Birdnest Jellyskin Lichen | | | | S2S3 | 8 | 22.5 ± 0.0 | NS |
| N | <i>Melanohalea septentrionalis</i> | Northern Camouflage Lichen | | | | S2S3 | 1 | 54.7 ± 0.0 | NS |
| N | <i>Myelochroa aurulenta</i> | Powdery Axil-bristle Lichen | | | | S2S3 | 4 | 22.7 ± 0.0 | NS |
| N | <i>Parmelia fertilis</i> | Fertile Shield Lichen | | | | S2S3 | 10 | 17.5 ± 0.0 | NS |
| N | <i>Hypotrachyna minarum</i> | Hairless-spined Shield Lichen | | | | S2S3 | 1 | 74.9 ± 0.0 | NS |
| N | <i>Parmeliopsis ambigua</i> | Green Starburst Lichen | | | | S2S3 | 3 | 72.3 ± 1.0 | NS |
| N | <i>Racodium rupestre</i> | Rockhair Lichen | | | | S2S3 | 1 | 69.9 ± 1.0 | NS |
| N | <i>Umbilicaria polyphylla</i> | Petalled Rocktripe Lichen | | | | S2S3 | 1 | 77.5 ± 2.0 | NS |
| N | <i>Usnea cavernosa</i> | Pitted Beard Lichen | | | | S2S3 | 2 | 55.2 ± 0.0 | NS |
| N | <i>Usnea mutabilis</i> | Bloody Beard Lichen | | | | S2S3 | 1 | 55.3 ± 0.0 | NS |
| N | <i>Fuscopannaria soreliata</i> | a Lichen | | | | S2S3 | 4 | 69.9 ± 1.0 | NS |
| N | <i>Stereocaulon condensatum</i> | Granular Soil Foam Lichen | | | | S2S3 | 6 | 57.2 ± 0.0 | NS |
| N | <i>Physcia subtilis</i> | Slender Rosette Lichen | | | | S2S3 | 1 | 62.8 ± 0.0 | NS |
| N | <i>Dimelaena oreina</i> | Golden Moonglow Lichen | | | | S2S3 | 1 | 90.9 ± 0.0 | NS |
| N | <i>Cladonia coccifera</i> | Eastern Boreal Pixie-cup Lichen | | | | S2S3 | 3 | 46.7 ± 4.0 | NS |
| N | <i>Cladonia deformis</i> | Lesser Sulphur-cup Lichen | | | | S2S3 | 4 | 71.7 ± 2.0 | NS |
| N | <i>Cladonia phyllophora</i> | Felt Lichen | | | | S2S3 | 2 | 86.7 ± 4.0 | NS |

| Taxonomic Group | Scientific Name | Common Name | COSEWIC | SARA | Prov Legal Prot | Prov Rarity Rank | # recs | Distance (km) | Prov |
|-----------------|----------------------------------|--------------------------------|---------|------|-----------------|------------------|--------|---------------|------|
| N | <i>Ephemerum serratum</i> | a Moss | | | | S3 | 6 | 25.9 ± 0.0 | NS |
| N | <i>Fissidens taxifolius</i> | Yew-leaved Pocket Moss | | | | S3 | 9 | 25.9 ± 0.0 | NS |
| N | <i>Anomodon tristis</i> | a Moss | | | | S3 | 9 | 64.7 ± 0.0 | NS |
| N | <i>Sphagnum contortum</i> | Twisted Peat Moss | | | | S3 | 4 | 35.3 ± 4.0 | NS |
| N | <i>Tetraplodon angustatus</i> | Toothed-leaved Nitrogen Moss | | | | S3 | 3 | 77.5 ± 0.0 | NS |
| N | <i>Rostania occultata</i> | Crusted Tarpaper Lichen | | | | S3 | 3 | 54.1 ± 0.0 | NS |
| N | <i>Collema nigrescens</i> | Blistered Tarpaper Lichen | | | | S3 | 29 | 35.1 ± 0.0 | NS |
| N | <i>Solorina saccata</i> | Woodland Owl Lichen | | | | S3 | 11 | 30.8 ± 4.0 | NS |
| N | <i>Fuscopannaria ahlneri</i> | Corrugated Shingles Lichen | | | | S3 | 73 | 20.5 ± 0.0 | NS |
| N | <i>Scytinium lichenoides</i> | Tattered Jellyskin Lichen | | | | S3 | 30 | 16.3 ± 0.0 | NS |
| N | <i>Leptogium milligranum</i> | Stretched Jellyskin Lichen | | | | S3 | 13 | 24.6 ± 3.0 | NS |
| N | <i>Nephroma bellum</i> | Naked Kidney Lichen | | | | S3 | 9 | 12.8 ± 0.0 | NS |
| N | <i>Placynthium nigrum</i> | Common Ink Lichen | | | | S3 | 2 | 56.3 ± 0.0 | NS |
| N | <i>Platismatia norvegica</i> | Oldgrowth Rag Lichen | | | | S3 | 1 | 91.9 ± 0.0 | NS |
| N | <i>Punctelia appalachensis</i> | Appalachian Speckleback Lichen | | | | S3 | 108 | 62.1 ± 0.0 | NS |
| N | <i>Viridothelium virens</i> | | | | | S3 | 2 | 64.5 ± 0.0 | NS |
| N | <i>Epebe lanata</i> | Waterside Rockshag Lichen | | | | S3 | 3 | 57.6 ± 17.0 | NS |
| N | <i>Phaeophyscia adlastola</i> | Powder-tipped Shadow Lichen | | | | S3 | 1 | 83.5 ± 0.0 | NS |
| N | <i>Phaeophyscia pusilloides</i> | Pompom-tipped Shadow Lichen | | | | S3 | 13 | 55.3 ± 0.0 | NS |
| N | <i>Peltigera collina</i> | Tree Pelt Lichen | | | | S3 | 6 | 26.8 ± 0.0 | NS |
| N | <i>Metzgeria conjugata</i> | Rock Veilwort | | | | S3? | 1 | 87.1 ± 0.0 | NS |
| N | <i>Barbula convoluta</i> | Lesser Bird's-claw Beard Moss | | | | S3? | 3 | 63.1 ± 0.0 | NS |
| N | <i>Calliergon giganteum</i> | Giant Spear Moss | | | | S3? | 2 | 58.0 ± 3.0 | NS |
| N | <i>Drummondia prorepens</i> | a Moss | | | | S3? | 1 | 72.0 ± 5.0 | NS |
| N | <i>Elodium blandowii</i> | Blandow's Bog Moss | | | | S3? | 6 | 51.5 ± 3.0 | NS |
| N | <i>Mnium stellare</i> | Star Leafy Moss | | | | S3? | 3 | 38.1 ± 1.0 | NS |
| N | <i>Sphagnum lindbergii</i> | Lindberg's Peat Moss | | | | S3? | 1 | 95.2 ± 0.0 | NS |
| N | <i>Sphagnum riparium</i> | Streamside Peat Moss | | | | S3? | 3 | 70.4 ± 0.0 | NS |
| N | <i>Cladonia stygia</i> | Black-footed Reindeer Lichen | | | | S3? | 17 | 67.8 ± 0.0 | NS |
| N | <i>Anomodon rugelii</i> | Rugel's Anomodon Moss | | | | S3S4 | 2 | 64.7 ± 0.0 | NS |
| N | <i>Dichelyma capillaceum</i> | Hairlike Dichelyma Moss | | | | S3S4 | 1 | 67.7 ± 3.0 | NS |
| N | <i>Encalypta ciliata</i> | Fringed Extinguisher Moss | | | | S3S4 | 2 | 62.0 ± 3.0 | NS |
| N | <i>Encalypta procera</i> | Slender Extinguisher Moss | | | | S3S4 | 1 | 63.9 ± 3.0 | NS |
| N | <i>Myurella julacea</i> | Small Mouse-tail Moss | | | | S3S4 | 1 | 35.8 ± 0.0 | NS |
| N | <i>Splachnum ampullaceum</i> | Cruet Dung Moss | | | | S3S4 | 4 | 71.3 ± 0.0 | NS |
| N | <i>Thamnobryum alleghaniense</i> | a Moss | | | | S3S4 | 8 | 33.0 ± 0.0 | NS |
| N | <i>Tomentypnum nitens</i> | Golden Fuzzy Fen Moss | | | | S3S4 | 2 | 53.5 ± 0.0 | NS |
| N | <i>Schistidium agassizii</i> | Elf Bloom Moss | | | | S3S4 | 1 | 78.4 ± 0.0 | NS |
| N | <i>Hylocomiastrum pyrenaicum</i> | a Feather Moss | | | | S3S4 | 2 | 63.9 ± 3.0 | NS |
| N | <i>Bryoria pseudofuscescens</i> | Mountain Horsehair Lichen | | | | S3S4 | 11 | 96.3 ± 0.0 | PE |
| N | <i>Enchylium tenax</i> | Soil Tarpaper Lichen | | | | S3S4 | 9 | 16.3 ± 0.0 | NS |
| N | <i>Sticta fuliginosa</i> | Peppered Moon Lichen | | | | S3S4 | 60 | 13.3 ± 1.0 | NS |
| N | <i>Arctoparmelia incurva</i> | Finger Ring Lichen | | | | S3S4 | 63 | 81.9 ± 0.0 | NS |
| N | <i>Scytinium teretiusculum</i> | Curly Jellyskin Lichen | | | | S3S4 | 14 | 44.3 ± 0.0 | NS |
| N | <i>Leptogium acadense</i> | Acadian Jellyskin Lichen | | | | S3S4 | 25 | 10.8 ± 0.0 | NS |
| N | <i>Scytinium subtile</i> | Appressed Jellyskin Lichen | | | | S3S4 | 29 | 30.8 ± 4.0 | NS |
| N | <i>Cladonia floerkeana</i> | Gritty British Soldiers Lichen | | | | S3S4 | 3 | 90.8 ± 0.0 | NS |
| N | <i>Vahlia leucophaea</i> | Shelter Shingle Lichen | | | | S3S4 | 10 | 32.1 ± 0.0 | NS |
| N | <i>Heterodermia speciosa</i> | Powdered Fringe Lichen | | | | S3S4 | 44 | 14.2 ± 0.0 | NS |
| N | <i>Leptogium corticola</i> | Blistered Jellyskin Lichen | | | | S3S4 | 67 | 51.1 ± 4.0 | NS |
| N | <i>Melanohalea olivacea</i> | Spotted Camouflage Lichen | | | | S3S4 | 6 | 55.2 ± 0.0 | NS |
| N | <i>Parmeliopsis hyperopta</i> | Gray Starburst Lichen | | | | S3S4 | 3 | 25.9 ± 0.0 | NS |

| Taxonomic Group | Scientific Name | Common Name | COSEWIC | SARA | Prov Legal Prot | Prov Rarity Rank | # recs | Distance (km) | Prov |
|-----------------|---|------------------------------------|-----------------|-----------------|-----------------|------------------|--------|---------------|------|
| N | <i>Parmotrema perlatum</i> | Powdered Ruffle Lichen | | | | S3S4 | 11 | 68.3 ± 0.0 | NS |
| N | <i>Peltigera hymenina</i> | Cloudy Pelt Lichen | | | | S3S4 | 2 | 94.1 ± 1.0 | NS |
| N | <i>Sphaerophorus fragilis</i> | Fragile Coral Lichen | | | | S3S4 | 3 | 83.1 ± 0.0 | NS |
| N | <i>Coccocarpia palmicola</i> | Salted Shell Lichen | | | | S3S4 | 524 | 40.3 ± 0.0 | NS |
| N | <i>Physcia caesia</i> | Blue-gray Rosette Lichen | | | | S3S4 | 1 | 75.8 ± 0.0 | NS |
| N | <i>Physcia tenella</i> | Fringed Rosette Lichen | | | | S3S4 | 5 | 78.8 ± 0.0 | NS |
| N | <i>Anaptychia palmulata</i> | Shaggy Fringed Lichen | | | | S3S4 | 104 | 22.7 ± 0.0 | NS |
| N | <i>Evernia prunastri</i> | Valley Oakmoss Lichen | | | | S3S4 | 37 | 20.1 ± 0.0 | NS |
| N | <i>Heterodermia neglecta</i> | Fringe Lichen | | | | S3S4 | 110 | 44.9 ± 0.0 | NS |
| P | <i>Clethra alnifolia</i> | Coast Pepper-Bush | Endangered | Threatened | Vulnerable | S2 | 2 | 86.7 ± 0.0 | NS |
| P | <i>Fraxinus nigra</i> | Black Ash | Threatened | | Threatened | S1S2 | 701 | 7.9 ± 1.0 | NS |
| P | <i>Bartonia paniculata</i> ssp. <i>paniculata</i> | Branched Bartonia | Threatened | Threatened | | SNA | 1 | 62.2 ± 10.0 | NS |
| P | <i>Lilaeopsis chinensis</i> | Eastern Lilaepsis | Special Concern | Special Concern | Vulnerable | S3 | 17 | 45.6 ± 1.0 | NS |
| P | <i>Isoetes prototypus</i> | Prototype Quillwort | Special Concern | Special Concern | Vulnerable | S3 | 13 | 16.6 ± 0.0 | NS |
| P | <i>Floerkea proserpinacoides</i> | False Mermaidweed | Not At Risk | | | S2S3 | 38 | 28.3 ± 7.0 | NS |
| P | <i>Acer saccharinum</i> | Silver Maple | | | | S1 | 11 | 83.9 ± 0.0 | NS |
| P | <i>Osmorhiza depauperata</i> | Blunt Sweet Cicely | | | | S1 | 1 | 72.4 ± 5.0 | NS |
| P | <i>Antennaria rosea</i> ssp. <i>arida</i> | Rosy Pussytoes | | | | S1 | 1 | 96.4 ± 0.0 | NS |
| P | <i>Andersonglossum boreale</i> | Northern Wild Comfrey | | | | S1 | 5 | 59.2 ± 1.0 | NS |
| P | <i>Lobelia spicata</i> | Pale-Spiked Lobelia | | | | S1 | 13 | 2.6 ± 7.0 | NS |
| P | <i>Hudsonia tomentosa</i> | Woolly Beach-heath | | | | S1 | 9 | 83.1 ± 7.0 | NS |
| P | <i>Callitriche hermaphroditica</i> | Northern Water-starwort | | | | S1 | 1 | 85.0 ± 0.0 | NB |
| P | <i>Elatine americana</i> | American Waterwort | | | | S1 | 2 | 56.0 ± 0.0 | NS |
| P | <i>Astragalus robbinsii</i> var. <i>minor</i> | Robbins' Milkvetch | | | | S1 | 28 | 96.3 ± 0.0 | NS |
| P | <i>Ribes americanum</i> | Wild Black Currant | | | | S1 | 4 | 17.0 ± 1.0 | NS |
| P | <i>Utricularia ochroleuca</i> | Yellowish-white Bladderwort | | | | S1 | 37 | 74.3 ± 0.0 | NS |
| P | <i>Fraxinus pennsylvanica</i> | Red Ash | | | | S1 | 13 | 43.9 ± 0.0 | NS |
| P | <i>Polygonum achoreum</i> | Leathery Knotweed | | | | S1 | 2 | 83.6 ± 0.0 | NB |
| P | <i>Persicaria careyi</i> | Carey's Smartweed | | | | S1 | 1 | 35.3 ± 3.0 | NS |
| P | <i>Phytolacca americana</i> | Common Pokeweed | | | | S1 | 1 | 76.7 ± 0.0 | NS |
| P | <i>Montia fontana</i> | Water Blinks | | | | S1 | 3 | 84.9 ± 1.0 | NS |
| P | <i>Lysimachia quadrifolia</i> | Whorled Yellow Loosestrife | | | | S1 | 1 | 80.9 ± 0.0 | NS |
| P | <i>Clematis occidentalis</i> | Purple Clematis | | | | S1 | 3 | 69.8 ± 0.0 | NS |
| P | <i>Ranunculus pensylvanicus</i> | Pennsylvania Buttercup | | | | S1 | 31 | 29.9 ± 0.0 | NS |
| P | <i>Amelanchier nantucketensis</i> | Nantucket Serviceberry | | | | S1 | 1 | 59.0 ± 1.0 | NS |
| P | <i>Salix myrtilifolia</i> | Blueberry Willow | | | | S1 | 1 | 52.3 ± 0.0 | NS |
| P | <i>Salix serissima</i> | Autumn Willow | | | | S1 | 2 | 52.3 ± 0.0 | NS |
| P | <i>Carex garberi</i> | Garber's Sedge | | | | S1 | 4 | 32.1 ± 0.0 | NS |
| P | <i>Carex granularis</i> | Limestone Meadow Sedge | | | | S1 | 2 | 77.5 ± 0.0 | NS |
| P | <i>Carex laxiflora</i> | Loose-Flowered Sedge | | | | S1 | 1 | 67.5 ± 1.0 | NS |
| P | <i>Carex ormostachya</i> | Necklace Spike Sedge | | | | S1 | 1 | 87.2 ± 1.0 | NB |
| P | <i>Carex plantaginea</i> | Plantain-Leaved Sedge | | | | S1 | 4 | 24.2 ± 0.0 | NS |
| P | <i>Carex prairea</i> | Prairie Sedge | | | | S1 | 2 | 82.2 ± 1.0 | NS |
| P | <i>Carex tenuiflora</i> | Sparse-Flowered Sedge | | | | S1 | 2 | 75.9 ± 0.0 | NS |
| P | <i>Carex viridula</i> var. <i>saxillitoralis</i> | Greenish Sedge | | | | S1 | 5 | 75.9 ± 0.0 | NS |
| P | <i>Cyperus lupulinus</i> ssp. <i>macilentus</i> | Hop Flatsedge | | | | S1 | 2 | 76.6 ± 0.0 | NS |
| P | <i>Scirpus atrovirens</i> | Dark-green Bulrush | | | | S1 | 4 | 43.1 ± 0.0 | NS |
| P | <i>Iris prismatica</i> | Slender Blue Flag | | | | S1 | 1 | 93.0 ± 100.0 | NS |
| P | <i>Juncus secundus</i> | Secund Rush | | | | S1 | 1 | 92.0 ± 0.0 | NS |
| P | <i>Juncus vaseyi</i> | Vasey Rush | | | | S1 | 4 | 34.3 ± 0.0 | NS |
| P | <i>Trillium grandiflorum</i> | White Trillium | | | | S1 | 3 | 82.2 ± 1.0 | NS |
| P | <i>Malaxis monophyllos</i> var. <i>brachypoda</i> | North American White Adder's-mouth | | | | S1 | 5 | 38.2 ± 1.0 | NS |
| P | <i>Spiranthes casei</i> var. <i>casei</i> | Case's Ladies'-Tresses | | | | S1 | 1 | 87.6 ± 0.0 | NS |

| Taxonomic Group | Scientific Name | Common Name | COSEWIC | SARA | Prov Legal Prot | Prov Rarity Rank | # recs | Distance (km) | Prov |
|-----------------|--|----------------------------|---------|------|-----------------|------------------|--------|---------------|------|
| P | <i>Calamagrostis stricta</i> ssp. <i>inexpansa</i> | Slim-stemmed Reed Grass | | | | S1 | 1 | 83.2 ± 1.0 | NB |
| P | <i>Elymus hystrix</i> | Spreading Wild Rye | | | | S1 | 12 | 16.5 ± 1.0 | NS |
| P | <i>Adiantum pedatum</i> | Northern Maidenhair Fern | | | | S1 | 14 | 19.3 ± 1.0 | NS |
| P | <i>Equisetum palustre</i> | Marsh Horsetail | | | | S1 | 1 | 82.5 ± 5.0 | NS |
| P | <i>Botrychium lunaria</i> | Common Moonwort | | | | S1 | 8 | 84.4 ± 0.0 | NS |
| P | <i>Selaginella rupestris</i> | Rock Spikemoss | | | | S1 | 1 | 59.1 ± 0.0 | NS |
| P | <i>Solidago hispida</i> | Hairy Goldenrod | | | | S1? | 2 | 81.2 ± 7.0 | NS |
| P | <i>Suaeda rolandii</i> | Roland's Sea-Blite | | | | S1? | 7 | 28.2 ± 2.0 | NS |
| P | <i>Carex pensylvanica</i> | Pennsylvania Sedge | | | | S1? | 3 | 29.9 ± 3.0 | NS |
| P | <i>Carex rostrata</i> | Narrow-leaved Beaked Sedge | | | | S1? | 1 | 79.1 ± 0.0 | NS |
| P | <i>Bolboschoenus robustus</i> | Sturdy Bulrush | | | | S1? | 2 | 47.2 ± 5.0 | NS |
| P | <i>Allium schoenoprasum</i> | Wild Chives | | | | S1? | 9 | 8.4 ± 0.0 | NS |
| P | <i>Allium schoenoprasum</i> var. <i>sibiricum</i> | Wild Chives | | | | S1? | 1 | 18.3 ± 7.0 | NS |
| P | <i>Crocanthemum canadense</i> | Long-branched Frostweed | | | Endangered | S1S2 | 2 | 83.3 ± 1.0 | NS |
| P | <i>Cypripedium arietinum</i> | Ram's-Head Lady's-Slipper | | | Endangered | S1S2 | 300 | 38.6 ± 0.0 | NS |
| P | <i>Sanicula odorata</i> | Clustered Sanicle | | | | S1S2 | 13 | 16.6 ± 10.0 | NS |
| P | <i>Ageratina altissima</i> | White Snakeroot | | | | S1S2 | 4 | 91.8 ± 10.0 | NS |
| P | <i>Draba glabella</i> | Rock Whitlow-Grass | | | | S1S2 | 4 | 62.1 ± 0.0 | NS |
| P | <i>Proserpinaca intermedia</i> | Intermediate Mermaidweed | | | | S1S2 | 2 | 48.3 ± 0.0 | NS |
| P | <i>Anemone virginiana</i> var. <i>alba</i> | Virginia Anemone | | | | S1S2 | 5 | 18.3 ± 7.0 | NS |
| P | <i>Carex haydenii</i> | Hayden's Sedge | | | | S1S2 | 3 | 20.2 ± 1.0 | NS |
| P | <i>Platanthera huronensis</i> | Fragrant Green Orchid | | | | S1S2 | 1 | 52.5 ± 10.0 | NS |
| P | <i>Calamagrostis stricta</i> ssp. <i>stricta</i> | Slim-stemmed Reed Grass | | | | S1S2 | 7 | 54.9 ± 7.0 | NS |
| P | <i>Carex vacillans</i> | Estuarine Sedge | | | | S1S3 | 5 | 76.1 ± 0.0 | NB |
| P | <i>Zizia aurea</i> | Golden Alexanders | | | | S2 | 38 | 28.2 ± 1.0 | NS |
| P | <i>Antennaria parlinii</i> ssp. <i>fallax</i> | Parlin's Pussytoes | | | | S2 | 13 | 33.9 ± 7.0 | NS |
| P | <i>Rudbeckia laciniata</i> | Cut-Leaved Coneflower | | | | S2 | 28 | 36.9 ± 0.0 | NS |
| P | <i>Rudbeckia laciniata</i> var. <i>laciniata</i> | Cut-Leaved Coneflower | | | | S2 | 7 | 72.6 ± 0.0 | NS |
| P | <i>Arabis pycnocarpa</i> | Cream-flowered Rockcress | | | | S2 | 1 | 41.8 ± 0.0 | NS |
| P | <i>Cardamine maxima</i> | Large Toothwort | | | | S2 | 2 | 70.9 ± 0.0 | NS |
| P | <i>Hudsonia ericoides</i> | Pinebarren Golden Heather | | | | S2 | 111 | 81.8 ± 7.0 | NS |
| P | <i>Desmodium canadense</i> | Canada Tick-trefoil | | | | S2 | 22 | 22.0 ± 5.0 | NS |
| P | <i>Hylodesmum glutinosum</i> | Large Tick-trefoil | | | | S2 | 21 | 49.9 ± 0.0 | NS |
| P | <i>Oxytropis campestris</i> var. <i>johannensis</i> | Field Locoweed | | | | S2 | 26 | 96.2 ± 1.0 | NS |
| P | <i>Conopholis americana</i> | American Cancer-root | | | | S2 | 3 | 79.2 ± 0.0 | NS |
| P | <i>Anemonastrum canadense</i> | Canada Anemone | | | | S2 | 8 | 22.9 ± 0.0 | NS |
| P | <i>Hepatica americana</i> | Round-lobed Hepatica | | | | S2 | 69 | 24.8 ± 7.0 | NS |
| P | <i>Ranunculus sceleratus</i> | Cursed Buttercup | | | | S2 | 22 | 71.6 ± 0.0 | NS |
| P | <i>Galium boreale</i> | Northern Bedstraw | | | | S2 | 10 | 23.1 ± 5.0 | NS |
| P | <i>Comandra umbellata</i> | Bastard's Toadflax | | | | S2 | 1 | 97.4 ± 1.0 | NB |
| P | <i>Gratiola neglecta</i> | Clammy Hedge-Hyssop | | | | S2 | 19 | 26.4 ± 2.0 | NS |
| P | <i>Dirca palustris</i> | Eastern Leatherwood | | | | S2 | 66 | 31.9 ± 7.0 | NS |
| P | <i>Carex chordorrhiza</i> | Creeping Sedge | | | | S2 | 52 | 72.5 ± 1.0 | NS |
| P | <i>Carex gynocrates</i> | Northern Bog Sedge | | | | S2 | 2 | 52.3 ± 0.0 | NS |
| P | <i>Carex pellita</i> | Woolly Sedge | | | | S2 | 12 | 30.9 ± 0.0 | NS |
| P | <i>Carex livida</i> | Livid Sedge | | | | S2 | 49 | 55.7 ± 0.0 | NS |
| P | <i>Juncus greenii</i> | Greene's Rush | | | | S2 | 11 | 17.1 ± 1.0 | NS |
| P | <i>Allium tricoccum</i> | Wild Leek | | | | S2 | 64 | 38.2 ± 0.0 | NS |
| P | <i>Lilium canadense</i> | Canada Lily | | | | S2 | 118 | 8.4 ± 7.0 | NS |
| P | <i>Cypripedium parviflorum</i> var. <i>pubescens</i> | Yellow Lady's-slipper | | | | S2 | 23 | 35.5 ± 0.0 | NS |

| Taxonomic Group | Scientific Name | Common Name | COSEWIC | SARA | Prov Legal Prot | Prov Rarity Rank | # recs | Distance (km) | Prov |
|-----------------|--|--------------------------------------|---------|------|-----------------|------------------|--------|---------------|------|
| P | <i>Cypripedium parviflorum</i> var. <i>makasin</i> | Small Yellow Lady's-Slipper | | | | S2 | 13 | 58.3 ± 0.0 | NS |
| P | <i>Cypripedium reginae</i> | Showy Lady's-Slipper | | | | S2 | 52 | 28.4 ± 7.0 | NS |
| P | <i>Platanthera flava</i> var. <i>flava</i> | Southern Rein Orchid | | | | S2 | 1 | 69.5 ± 7.0 | NS |
| P | <i>Platanthera flava</i> var. <i>herbiola</i> | Pale Green Orchid | | | | S2 | 11 | 37.8 ± 1.0 | NS |
| P | <i>Platanthera macrophylla</i> | Large Round-Leaved Orchid | | | | S2 | 15 | 9.6 ± 1.0 | NS |
| P | <i>Bromus latiglumis</i> | Broad-Glumed Brome | | | | S2 | 33 | 38.6 ± 0.0 | NS |
| P | <i>Cinna arundinacea</i> | Sweet Wood Reed Grass | | | | S2 | 20 | 38.8 ± 0.0 | NS |
| P | <i>Elymus wiegandii</i> | Wiegand's Wild Rye | | | | S2 | 21 | 31.5 ± 0.0 | NS |
| P | <i>Festuca subverticillata</i> | Nodding Fescue | | | | S2 | 13 | 19.3 ± 1.0 | NS |
| P | <i>Cryptogramma stelleri</i> | Steller's Rockbrake | | | | S2 | 3 | 28.0 ± 0.0 | NS |
| P | <i>Cuscuta cephalanthi</i> | Buttonbush Dodder | | | | S2? | 3 | 69.0 ± 1.0 | NS |
| P | <i>Rumex persicarioides</i> | Peach-leaved Dock | | | | S2? | 1 | 96.3 ± 5.0 | PE |
| P | <i>Crataegus submollis</i> | Quebec Hawthorn | | | | S2? | 5 | 23.3 ± 7.0 | NS |
| P | <i>Carex peckii</i> | White-Tinged Sedge | | | | S2? | 4 | 16.8 ± 0.0 | NS |
| P | <i>Thuja occidentalis</i> | Eastern White Cedar | | | Vulnerable | S2S3 | 965 | 38.7 ± 0.0 | NS |
| P | <i>Osmorhiza longistylis</i> | Smooth Sweet Cicely | | | | S2S3 | 33 | 22.3 ± 5.0 | NS |
| P | <i>Bidens hyperborea</i> | Estuary Beggarticks | | | | S2S3 | 2 | 46.3 ± 0.0 | NS |
| P | <i>Erigeron philadelphicus</i> | Philadelphia Fleabane | | | | S2S3 | 5 | 55.4 ± 1.0 | NS |
| P | <i>Lactuca hirsuta</i> | Hairy Lettuce | | | | S2S3 | 4 | 72.2 ± 2.0 | NS |
| P | <i>Impatiens pallida</i> | Pale Jewelweed | | | | S2S3 | 4 | 26.4 ± 0.0 | NS |
| P | <i>Caulophyllum thalictroides</i> | Blue Cohosh | | | | S2S3 | 95 | 12.0 ± 7.0 | NS |
| P | <i>Draba arabisans</i> | Rock Whitlow-Grass | | | | S2S3 | 19 | 56.5 ± 0.0 | NS |
| P | <i>Boechera stricta</i> | Drummond's Rockcress | | | | S2S3 | 12 | 28.4 ± 0.0 | NS |
| P | <i>Stellaria humifusa</i> | Saltmarsh Starwort | | | | S2S3 | 11 | 38.9 ± 1.0 | NS |
| P | <i>Oxybasis rubra</i> | Red Goosefoot | | | | S2S3 | 4 | 73.2 ± 0.0 | NS |
| P | <i>Hypericum majus</i> | Large St John's-wort | | | | S2S3 | 20 | 33.9 ± 0.0 | NS |
| P | <i>Hypericum x dissimulatum</i> | Disguised St. John's-wort | | | | S2S3 | 8 | 20.9 ± 1.0 | NS |
| P | <i>Empetrum atropurpureum</i> | Purple Crowberry | | | | S2S3 | 4 | 81.6 ± 7.0 | NS |
| P | <i>Euphorbia polygonifolia</i> | Seaside Spurge | | | | S2S3 | 4 | 79.3 ± 0.0 | NB |
| P | <i>Myriophyllum farwellii</i> | Farwell's Water Milfoil | | | | S2S3 | 13 | 34.0 ± 1.0 | NS |
| P | <i>Hedeoma pulegioides</i> | American False Pennyroyal | | | | S2S3 | 17 | 40.9 ± 1.0 | NS |
| P | <i>Oenothera fruticosa</i> ssp. <i>tetragona</i> | Narrow-leaved Evening Primrose | | | | S2S3 | 7 | 18.3 ± 7.0 | NS |
| P | <i>Polygala polygama</i> | Racemed Milkwort | | | | S2S3 | 1 | 82.9 ± 1.0 | NS |
| P | <i>Polygonum aviculare</i> ssp. <i>buxiforme</i> | Box Knotweed | | | | S2S3 | 7 | 12.4 ± 0.0 | NS |
| P | <i>Polygonum oxyspermum</i> | Sharp-fruit Knotweed | | | | S2S3 | 1 | 70.8 ± 0.0 | NS |
| P | <i>Rumex triangulivalvis</i> | Triangular-valve Dock | | | | S2S3 | 12 | 10.3 ± 0.0 | NS |
| P | <i>Primula mistassinica</i> | Mistassini Primrose | | | | S2S3 | 16 | 18.3 ± 7.0 | NS |
| P | <i>Anemone quinquefolia</i> | Wood Anemone | | | | S2S3 | 18 | 30.6 ± 0.0 | NS |
| P | <i>Caltha palustris</i> | Yellow Marsh Marigold | | | | S2S3 | 15 | 71.7 ± 2.0 | NS |
| P | <i>Amelanchier fernaldii</i> | Fernald's Serviceberry | | | | S2S3 | 1 | 55.3 ± 5.0 | NS |
| P | <i>Potentilla canadensis</i> | Canada Cinquefoil | | | | S2S3 | 5 | 17.1 ± 5.0 | NS |
| P | <i>Galium obtusum</i> | Blunt-leaved Bedstraw | | | | S2S3 | 1 | 87.2 ± 1.0 | NB |
| P | <i>Salix pellita</i> | Satiny Willow | | | | S2S3 | 8 | 34.0 ± 4.0 | NS |
| P | <i>Tiarella cordifolia</i> | Heart-leaved Foamflower | | | | S2S3 | 232 | 19.3 ± 1.0 | NS |
| P | <i>Agalinis purpurea</i> var. <i>parviflora</i> | Small-flowered Purple False Foxglove | | | | S2S3 | 25 | 40.8 ± 0.0 | NS |
| P | <i>Boehmeria cylindrica</i> | Small-spike False-nettle | | | | S2S3 | 2 | 44.5 ± 0.0 | NS |
| P | <i>Carex adusta</i> | Lesser Brown Sedge | | | | S2S3 | 9 | 11.6 ± 7.0 | NS |
| P | <i>Carex capillaris</i> | Hairlike Sedge | | | | S2S3 | 9 | 45.0 ± 0.0 | NS |
| P | <i>Carex comosa</i> | Bearded Sedge | | | | S2S3 | 11 | 9.7 ± 0.0 | NS |
| P | <i>Carex houghtoniana</i> | Houghton's Sedge | | | | S2S3 | 5 | 29.9 ± 1.0 | NS |
| P | <i>Carex hystericina</i> | Porcupine Sedge | | | | S2S3 | 9 | 29.7 ± 1.0 | NS |
| P | <i>Eleocharis ovata</i> | Ovate Spikerush | | | | S2S3 | 16 | 24.4 ± 0.0 | NS |
| P | <i>Scirpus pedicellatus</i> | Stalked Bulrush | | | | S2S3 | 7 | 39.6 ± 1.0 | NS |

| Taxonomic Group | Scientific Name | Common Name | COSEWIC | SARA | Prov Legal Prot | Prov Rarity Rank | # recs | Distance (km) | Prov |
|-----------------|--|-------------------------------|---------|------|-----------------|------------------|--------|---------------|------|
| P | <i>Vallisneria americana</i> | Wild Celery | | | | S2S3 | 9 | 26.7 ± 1.0 | NS |
| P | <i>Najas gracillima</i> | Thread-Like Naiad | | | | S2S3 | 2 | 58.6 ± 0.0 | NS |
| P | <i>Goodyera pubescens</i> | Downy Rattlesnake-Plantain | | | | S2S3 | 14 | 41.2 ± 0.0 | NS |
| P | <i>Spiranthes lucida</i> | Shining Ladies'-Tresses | | | | S2S3 | 28 | 30.5 ± 0.0 | NS |
| P | <i>Calamagrostis stricta</i> | Slim-stemmed Reed Grass | | | | S2S3 | 5 | 74.2 ± 0.0 | NS |
| P | <i>Potamogeton friesii</i> | Fries' Pondweed | | | | S2S3 | 15 | 24.4 ± 5.0 | NS |
| P | <i>Woodsia glabella</i> | Smooth Cliff Fern | | | | S2S3 | 2 | 12.2 ± 1.0 | NS |
| P | <i>Botrychium lanceolatum</i> ssp. <i>angustisegmentum</i> | Narrow Triangle Moonwort | | | | S2S3 | 16 | 29.3 ± 0.0 | NS |
| P | <i>Botrychium simplex</i> | Least Moonwort | | | | S2S3 | 6 | 29.3 ± 0.0 | NS |
| P | <i>Ophioglossum pusillum</i> | Northern Adder's-tongue | | | | S2S3 | 9 | 45.0 ± 0.0 | NS |
| P | <i>Potamogeton pulcher</i> | Spotted Pondweed | | | Vulnerable | S3 | 3 | 55.1 ± 2.0 | NS |
| P | <i>Angelica atropurpurea</i> | Purple-stemmed Angelica | | | | S3 | 3 | 41.9 ± 0.0 | NS |
| P | <i>Conioselinum chinense</i> | Chinese Hemlock-parsley | | | | S3 | 10 | 29.4 ± 0.0 | NS |
| P | <i>Hieracium robinsonii</i> | Robinson's Hawkweed | | | | S3 | 3 | 22.7 ± 1.0 | NS |
| P | <i>Iva frutescens</i> | Big-leaved Marsh-elder | | | | S3 | 35 | 56.2 ± 0.0 | NS |
| P | <i>Senecio pseudoarnica</i> | Seabeach Ragwort | | | | S3 | 19 | 18.3 ± 7.0 | NS |
| P | <i>Symphyotrichum boreale</i> | Boreal Aster | | | | S3 | 6 | 18.3 ± 7.0 | NS |
| P | <i>Symphyotrichum undulatum</i> | Wavy-leaved Aster | | | | S3 | 7 | 59.2 ± 0.0 | NS |
| P | <i>Symphyotrichum ciliolatum</i> | Fringed Blue Aster | | | | S3 | 24 | 16.8 ± 0.0 | NS |
| P | <i>Betula pumila</i> var. <i>pumila</i> | Bog Birch | | | | S3 | 1 | 68.1 ± 1.0 | NS |
| P | <i>Betula michauxii</i> | Michaux's Dwarf Birch | | | | S3 | 27 | 54.2 ± 0.0 | NS |
| P | <i>Betula pumila</i> | Bog Birch | | | | S3 | 3 | 52.8 ± 0.0 | NS |
| P | <i>Cardamine parviflora</i> | Small-flowered Bittercress | | | | S3 | 12 | 62.0 ± 0.0 | NS |
| P | <i>Palustricodon aparinoides</i> | Marsh Bellflower | | | | S3 | 39 | 18.3 ± 5.0 | NS |
| P | <i>Mononeuria groenlandica</i> | Greenland Stitchwort | | | | S3 | 61 | 61.0 ± 0.0 | NS |
| P | <i>Sagina nodosa</i> | Knotted Pearlwort | | | | S3 | 11 | 82.7 ± 1.0 | NS |
| P | <i>Sagina nodosa</i> ssp. <i>borealis</i> | Knotted Pearlwort | | | | S3 | 7 | 96.2 ± 0.0 | NS |
| P | <i>Stellaria longifolia</i> | Long-leaved Starwort | | | | S3 | 19 | 20.3 ± 0.0 | NS |
| P | <i>Ceratophyllum echinatum</i> | Prickly Hornwort | | | | S3 | 41 | 9.7 ± 0.0 | NS |
| P | <i>Triosteum aurantiacum</i> | Orange-fruited Tinker's Weed | | | | S3 | 66 | 22.5 ± 0.0 | NS |
| P | <i>Viburnum edule</i> | Squashberry | | | | S3 | 3 | 49.7 ± 0.0 | NS |
| P | <i>Empetrum eamesii</i> | Pink Crowberry | | | | S3 | 15 | 81.6 ± 7.0 | NS |
| P | <i>Vaccinium uliginosum</i> | Alpine Bilberry | | | | S3 | 1 | 97.6 ± 1.0 | NS |
| P | <i>Geranium bicknellii</i> | Bicknell's Crane's-bill | | | | S3 | 13 | 39.0 ± 2.0 | NS |
| P | <i>Myriophyllum verticillatum</i> | Whorled Water Milfoil | | | | S3 | 13 | 38.6 ± 7.0 | NS |
| P | <i>Epilobium strictum</i> | Downy Willowherb | | | | S3 | 28 | 22.0 ± 5.0 | NS |
| P | <i>Polygala sanguinea</i> | Blood Milkwort | | | | S3 | 30 | 23.3 ± 5.0 | NS |
| P | <i>Persicaria arifolia</i> | Halberd-leaved Tearthumb | | | | S3 | 81 | 28.9 ± 0.0 | NS |
| P | <i>Plantago rugelii</i> | Rugel's Plantain | | | | S3 | 11 | 16.8 ± 0.0 | NS |
| P | <i>Primula laurentiana</i> | Laurentian Primrose | | | | S3 | 24 | 68.2 ± 0.0 | NS |
| P | <i>Samolus parviflorus</i> | Seaside Brookweed | | | | S3 | 14 | 42.1 ± 0.0 | NS |
| P | <i>Pyrola minor</i> | Lesser Pyrola | | | | S3 | 4 | 45.3 ± 0.0 | NS |
| P | <i>Anemone virginiana</i> | Virginia Anemone | | | | S3 | 17 | 14.4 ± 7.0 | NS |
| P | <i>Cephalanthus occidentalis</i> | Common Buttonbush | | | | S3 | 6 | 85.2 ± 0.0 | NS |
| P | <i>Galium labradoricum</i> | Labrador Bedstraw | | | | S3 | 83 | 41.9 ± 0.0 | NS |
| P | <i>Salix pedicellaris</i> | Bog Willow | | | | S3 | 66 | 44.1 ± 0.0 | NS |
| P | <i>Salix sericea</i> | Silky Willow | | | | S3 | 1 | 51.9 ± 1.0 | NS |
| P | <i>Saxifraga paniculata</i> ssp. <i>laestadii</i> | Laestadius' Saxifrage | | | | S3 | 12 | 62.1 ± 1.0 | NS |
| P | <i>Lindernia dubia</i> | Yellow-seeded False Pimpernel | | | | S3 | 48 | 30.4 ± 0.0 | NS |
| P | <i>Laportea canadensis</i> | Canada Wood Nettle | | | | S3 | 59 | 12.3 ± 0.0 | NS |
| P | <i>Pilea pumila</i> | Dwarf Clearweed | | | | S3 | 8 | 23.6 ± 0.0 | NS |
| P | <i>Viola nephrophylla</i> | Northern Bog Violet | | | | S3 | 10 | 21.9 ± 7.0 | NS |
| P | <i>Carex bebbii</i> | Bebb's Sedge | | | | S3 | 30 | 30.6 ± 0.0 | NS |
| P | <i>Carex castanea</i> | Chestnut Sedge | | | | S3 | 26 | 40.2 ± 0.0 | NS |

| Taxonomic Group | Scientific Name | Common Name | COSEWIC | SARA | Prov Legal Prot | Prov Rarity Rank | # recs | Distance (km) | Prov |
|-----------------|---|-----------------------------|---------|------|-----------------|------------------|--------|---------------|------|
| P | <i>Carex cryptolepis</i> | Hidden-scaled Sedge | | | | S3 | 18 | 20.1 ± 1.0 | NS |
| P | <i>Carex eburnea</i> | Bristle-leaved Sedge | | | | S3 | 10 | 16.8 ± 0.0 | NS |
| P | <i>Carex hirtifolia</i> | Pubescent Sedge | | | | S3 | 54 | 12.7 ± 0.0 | NS |
| P | <i>Carex lupulina</i> | Hop Sedge | | | | S3 | 48 | 20.1 ± 1.0 | NS |
| P | <i>Carex rosea</i> | Rosy Sedge | | | | S3 | 43 | 16.9 ± 0.0 | NS |
| P | <i>Carex swanii</i> | Swan's Sedge | | | | S3 | 3 | 72.1 ± 0.0 | NS |
| P | <i>Carex tenera</i> | Tender Sedge | | | | S3 | 11 | 14.9 ± 0.0 | NS |
| P | <i>Carex tribuloides</i> | Blunt Broom Sedge | | | | S3 | 15 | 18.9 ± 0.0 | NS |
| P | <i>Carex tuckermanii</i> | Tuckerman's Sedge | | | | S3 | 48 | 27.4 ± 0.0 | NS |
| P | <i>Carex atratiformis</i> | Scabrous Black Sedge | | | | S3 | 3 | 55.0 ± 0.0 | NS |
| P | <i>Eleocharis nitida</i> | Quill Spikerush | | | | S3 | 16 | 32.8 ± 7.0 | NS |
| P | <i>Eleocharis flavescens</i> var. <i>olivacea</i> | Bright-green Spikerush | | | | S3 | 5 | 33.4 ± 0.0 | NS |
| P | <i>Eriophorum gracile</i> | Slender Cottongrass | | | | S3 | 54 | 22.1 ± 10.0 | NS |
| P | <i>Juncus stygius</i> ssp. <i>americanus</i> | Moor Rush | | | | S3 | 75 | 74.8 ± 0.0 | NS |
| P | <i>Coeloglossum viride</i> | Long-bracted Frog Orchid | | | | S3 | 13 | 15.4 ± 0.0 | NS |
| P | <i>Cypripedium parviflorum</i> | Yellow Lady's-slipper | | | | S3 | 551 | 35.6 ± 0.0 | NS |
| P | <i>Neottia bifolia</i> | Southern Twayblade | | | | S3 | 101 | 28.6 ± 0.0 | NS |
| P | <i>Platanthera grandiflora</i> | Large Purple Fringed Orchid | | | | S3 | 122 | 14.7 ± 0.0 | NS |
| P | <i>Platanthera hookeri</i> | Hooker's Orchid | | | | S3 | 27 | 38.8 ± 0.0 | NS |
| P | <i>Dichanthelium linearifolium</i> | Narrow-leaved Panic Grass | | | | S3 | 7 | 30.8 ± 0.0 | NS |
| P | <i>Piptatheropsis canadensis</i> | Canada Ricegrass | | | | S3 | 8 | 8.9 ± 0.0 | NS |
| P | <i>Poa glauca</i> | Glaucous Blue Grass | | | | S3 | 8 | 45.0 ± 0.0 | NS |
| P | <i>Potamogeton praelongus</i> | White-stemmed Pondweed | | | | S3 | 11 | 11.8 ± 5.0 | NS |
| P | <i>Potamogeton richardsonii</i> | Richardson's Pondweed | | | | S3 | 10 | 29.6 ± 0.0 | NS |
| P | <i>Potamogeton zosteriformis</i> | Flat-stemmed Pondweed | | | | S3 | 29 | 26.7 ± 2.0 | NS |
| P | <i>Asplenium viride</i> | Green Spleenwort | | | | S3 | 12 | 21.7 ± 7.0 | NS |
| P | <i>Dryopteris fragrans</i> | Fragrant Wood Fern | | | | S3 | 15 | 12.3 ± 1.0 | NS |
| P | <i>Sceptridium dissectum</i> | Dissected Moonwort | | | | S3 | 8 | 29.3 ± 0.0 | NS |
| P | <i>Polypodium appalachianum</i> | Appalachian Polypody | | | | S3 | 18 | 15.5 ± 0.0 | NS |
| P | <i>Persicaria amphibia</i> var. <i>emersa</i> | Long-root Smartweed | | | | S3? | 3 | 44.6 ± 0.0 | NS |
| P | <i>Spiranthes ochroleuca</i> | Yellow Ladies'-tresses | | | | S3? | 26 | 15.0 ± 0.0 | NS |
| P | <i>Diphasiastrum x sabinifolium</i> | Savin-leaved Ground-cedar | | | | S3? | 9 | 9.2 ± 0.0 | NS |
| P | <i>Bidens vulgata</i> | Tall Beggarticks | | | | S3S4 | 7 | 20.8 ± 0.0 | NS |
| P | <i>Erigeron hyssopifolius</i> | Hyssop-leaved Fleabane | | | | S3S4 | 24 | 6.9 ± 0.0 | NS |
| P | <i>Hieracium paniculatum</i> | Panicled Hawkweed | | | | S3S4 | 19 | 43.0 ± 0.0 | NS |
| P | <i>Bidens beckii</i> | Water Beggarticks | | | | S3S4 | 26 | 16.9 ± 0.0 | NS |
| P | <i>Packera paupercula</i> | Balsam Groundsel | | | | S3S4 | 94 | 16.3 ± 0.0 | NS |
| P | <i>Packera paupercula</i> var. <i>paupercula</i> | Balsam Groundsel | | | | S3S4 | 1 | 60.0 ± 0.0 | NS |
| P | <i>Atriplex glabriuscula</i> var. <i>franktonii</i> | Frankton's Saltbush | | | | S3S4 | 18 | 53.6 ± 5.0 | NS |
| P | <i>Shepherdia canadensis</i> | Soapberry | | | | S3S4 | 101 | 56.0 ± 0.0 | NS |
| P | <i>Vaccinium boreale</i> | Northern Blueberry | | | | S3S4 | 4 | 54.6 ± 1.0 | NS |
| P | <i>Vaccinium cespitosum</i> | Dwarf Bilberry | | | | S3S4 | 59 | 29.4 ± 0.0 | NS |
| P | <i>Vaccinium corymbosum</i> | Highbush Blueberry | | | | S3S4 | 3 | 70.2 ± 0.0 | NS |
| P | <i>Fagus grandifolia</i> | American Beech | | | | S3S4 | 320 | 7.9 ± 1.0 | NS |
| P | <i>Bartonia virginica</i> | Yellow Bartonia | | | | S3S4 | 1 | 51.9 ± 7.0 | NS |
| P | <i>Proserpinaca pectinata</i> | Comb-leaved Mermaidweed | | | | S3S4 | 5 | 60.9 ± 5.0 | NS |
| P | <i>Nuphar microphylla</i> | Small Yellow Pond-lily | | | | S3S4 | 9 | 31.0 ± 1.0 | NS |
| P | <i>Persicaria pensylvanica</i> | Pennsylvania Smartweed | | | | S3S4 | 38 | 18.3 ± 7.0 | NS |
| P | <i>Fallopia scandens</i> | Climbing False Buckwheat | | | | S3S4 | 35 | 18.3 ± 7.0 | NS |
| P | <i>Rumex pallidus</i> | Seabeach Dock | | | | S3S4 | 1 | 80.4 ± 0.0 | NS |
| P | <i>Pyrola asarifolia</i> | Pink Pyrola | | | | S3S4 | 12 | 19.4 ± 1.0 | NS |
| P | <i>Endotropis alnifolia</i> | alder-leaved buckthorn | | | | S3S4 | 219 | 25.5 ± 1.0 | NS |
| P | <i>Amelanchier spicata</i> | Running Serviceberry | | | | S3S4 | 18 | 37.1 ± 5.0 | NS |

| Taxonomic Group | Scientific Name | Common Name | COSEWIC | SARA | Prov Legal Prot | Prov Rarity Rank | # recs | Distance (km) | Prov |
|-----------------|--|-----------------------------|---------|------|-----------------|------------------|--------|---------------|------|
| P | <i>Crataegus succulenta</i> | Fleshy Hawthorn | | | | S3S4 | 1 | 72.3 ± 0.0 | NS |
| P | <i>Fragaria vesca</i> ssp. <i>americana</i> | Woodland Strawberry | | | | S3S4 | 71 | 18.6 ± 0.0 | NS |
| P | <i>Fragaria vesca</i> | Woodland Strawberry | | | | S3S4 | 2 | 46.5 ± 0.0 | NS |
| P | <i>Galium aparine</i> | Common Bedstraw | | | | S3S4 | 28 | 20.7 ± 0.0 | NS |
| P | <i>Geocaulon lividum</i> | Northern Comandra | | | | S3S4 | 11 | 41.0 ± 0.0 | NS |
| P | <i>Limosella australis</i> | Southern Mudwort | | | | S3S4 | 24 | 43.2 ± 0.0 | NS |
| P | <i>Ulmus americana</i> | White Elm | | | | S3S4 | 94 | 7.9 ± 1.0 | NS |
| P | <i>Verbena hastata</i> | Blue Vervain | | | | S3S4 | 218 | 16.6 ± 0.0 | NS |
| P | <i>Viola sagittata</i> var. <i>ovata</i> | Arrow-Leaved Violet | | | | S3S4 | 17 | 55.0 ± 0.0 | NS |
| P | <i>Viola selkirkii</i> | Great-Spurred Violet | | | | S3S4 | 9 | 15.7 ± 0.0 | NS |
| P | <i>Symplocarpus foetidus</i> | Eastern Skunk Cabbage | | | | S3S4 | 133 | 71.9 ± 0.0 | NS |
| P | <i>Carex argyrantha</i> | Silvery-flowered Sedge | | | | S3S4 | 8 | 62.0 ± 2.0 | NS |
| P | <i>Triglochin gaspensis</i> | Gasp Arrowgrass | | | | S3S4 | 19 | 81.6 ± 1.0 | NB |
| P | <i>Juncus acuminatus</i> | Sharp-Fruit Rush | | | | S3S4 | 7 | 45.0 ± 0.0 | NS |
| P | <i>Juncus subcaudatus</i> | Woods-Rush | | | | S3S4 | 25 | 0.5 ± 5.0 | NS |
| P | <i>Luzula parviflora</i> ssp. <i>melanocarpa</i> | Black-fruited Woodrush | | | | S3S4 | 5 | 26.9 ± 0.0 | NS |
| P | <i>Goodyera repens</i> | Lesser Rattlesnake-plantain | | | | S3S4 | 20 | 41.3 ± 0.0 | NS |
| P | <i>Liparis loeselii</i> | Loesel's Twayblade | | | | S3S4 | 16 | 16.4 ± 1.0 | NS |
| P | <i>Platanthera obtusata</i> | Blunt-leaved Orchid | | | | S3S4 | 9 | 20.3 ± 1.0 | NS |
| P | <i>Platanthera orbiculata</i> | Small Round-leaved Orchid | | | | S3S4 | 34 | 11.6 ± 7.0 | NS |
| P | <i>Alopecurus aequalis</i> | Short-awned Foxtail | | | | S3S4 | 28 | 23.3 ± 0.0 | NS |
| P | <i>Dichanthelium clandestinum</i> | Deer-tongue Panic Grass | | | | S3S4 | 107 | 34.8 ± 0.0 | NS |
| P | <i>Panicum philadelphicum</i> | Philadelphia Panicgrass | | | | S3S4 | 15 | 30.2 ± 0.0 | NS |
| P | <i>Koeleria spicata</i> | Narrow False Oats | | | | S3S4 | 21 | 23.2 ± 0.0 | NS |
| P | <i>Asplenium trichomanes</i> | Maidenhair Spleenwort | | | | S3S4 | 15 | 60.4 ± 1.0 | NS |
| P | <i>Equisetum pratense</i> | Meadow Horsetail | | | | S3S4 | 17 | 6.2 ± 0.0 | NS |
| P | <i>Diphasiastrum complanatum</i> | Northern Ground-cedar | | | | S3S4 | 20 | 24.4 ± 0.0 | NS |
| P | <i>Diphasiastrum sitchense</i> | Sitka Ground-cedar | | | | S3S4 | 6 | 24.6 ± 5.0 | NS |
| P | <i>Huperzia appressa</i> | Mountain Firmoss | | | | S3S4 | 18 | 2.6 ± 7.0 | NS |
| P | <i>Sceptridium multifidum</i> | Leathery Moonwort | | | | S3S4 | 13 | 23.3 ± 10.0 | NS |
| P | <i>Botrychium matricariifolium</i> | Daisy-leaved Moonwort | | | | S3S4 | 11 | 38.8 ± 1.0 | NS |
| P | <i>Viola canadensis</i> | Canada Violet | | | | SH | 2 | 28.3 ± 7.0 | NS |
| P | <i>Greeneochloa coarctata</i> | Small Reedgrass | | | | SH | 1 | 84.3 ± 6.0 | NS |

5.1 SOURCE BIBLIOGRAPHY (100 km)

The recipient of these data shall acknowledge the AC CDC and the data sources listed below in any documents, reports, publications or presentations, in which this dataset makes a significant contribution.

| # recs | CITATION |
|--------|---|
| 13546 | Morrison, Guy. 2011. Maritime Shorebird Survey (MSS) database. Canadian Wildlife Service, Ottawa, 15939 surveys. 86171 recs. |
| 10618 | Lepage, D. 2014. Maritime Breeding Bird Atlas Database. Bird Studies Canada, Sackville NB, 407,838 recs. |
| 4525 | Pardieck, K.L., Ziolkowski Jr., D.J., Lutmerding, M., Aponete, V.I., and Hudson, M-A.R. 2020. North American Breeding Bird Survey Dataset 1966 - 2019: U.S. Geological Survey data release, https://doi.org/10.5066/P9J6QUF6 |
| 4468 | Erskine, A.J. 1992. Maritime Breeding Bird Atlas Database. NS Museum & Nimbus Publ., Halifax, 82,125 recs. |
| 4021 | Paquet, Julie. 2018. Atlantic Canada Shorebird Survey (ACSS) database 2012-2018. Environment Canada, Canadian Wildlife Service. |
| 2534 | iNaturalist. 2020. iNaturalist Data Export 2020. iNaturalist.org and iNaturalist.ca, Web site: 128728 recs. |
| 2119 | Berrigan, L. 2019. Maritimes Marsh Monitoring Project 2013, 2014, 2016, 2017, and 2018 data. Bird Studies Canada, Sackville, NB. |
| 1930 | eBird. 2014. eBird Basic Dataset. Version: EBD_relNov-2014. Ithaca, New York. Nov 2014. Cornell Lab of Ornithology, 25036 recs. |
| 1859 | eBird. 2020. eBird Basic Dataset. Version: EBD_relNov-2019. Ithaca, New York. Nov 2019, Cape Breton Bras d'Or Lakes Watershed subset. Cornell Lab of Ornithology. |
| 923 | SwiftWatch. 2022. Total Chimney Swift counts from roost watches for the duration of the SwiftWatch program (2011-2021). Birds Canada. |
| 897 | Patrick, A.; Horne, D.; Noseworthy, J. et. al. 2017. Field data for Nova Scotia and New Brunswick, 2015 and 2017. Nature Conservancy of Canada. |
| 718 | Eaton, S. 2014. Nova Scotia Wood Turtle Database. Environment and Climate Change Canada, 4843 recs. |
| 622 | Cameron, E. 2008. Canadian Gypsum Co. survey 2007-08. Conestoga-Rovers & Assoc., 623 recs. |

| # recs | CITATION |
|--------|---|
| 569 | Blaney, C.S.; Mazerolle, D.M.; Belliveau, A.B. 2015. Atlantic Canada Conservation Data Centre Fieldwork 2015. Atlantic Canada Conservation Data Centre, # recs. |
| 488 | Neily, T.H. & Pepper, C.; Toms, B. 2018. Nova Scotia lichen database [as of 2018-03]. Mersey Tobeatic Research Institute. |
| 447 | Belliveau, A.G. 2020. E.C. Smith Herbarium and Atlantic Canada Conservation Data Centre Fieldwork 2019, 2020. E.C. Smith Herbarium. |
| 436 | Tranquilla, L. 2015. Maritimes Marsh Monitoring Project 2015 data. Bird Studies Canada, Sackville NB, 5062 recs. |
| 423 | Paquet, Julie. 2019. Atlantic Canada Shorebird Survey ACSG database for 2019. Environment Canada, Canadian Wildlife Service. |
| 421 | Clayden, S. Digitization of Wolfgang Maass Nova Scotia forest lichen collections, 1964-2004. New Brunswick Museum. 2018. |
| 418 | Benjamin, L.K. (compiler). 2007. Significant Habitat & Species Database. Nova Scotia Dept Natural Resources, 8439 recs. |
| 414 | Newell, R.E. 2000. E.C. Smith Herbarium Database. Acadia University, Wolfville NS, 7139 recs. |
| 400 | Amirault, D.L. & Stewart, J. 2007. Piping Plover Database 1894-2006. Canadian Wildlife Service, Sackville, 3344 recs, 1228 new. |
| 370 | Newell, R.E. 2005. E.C. Smith Digital Herbarium. E.C. Smith Herbarium, Irving Biodiversity Collection, Acadia University, Web site: http://luxor.acadiau.ca/library/Herbarium/project/ . 582 recs. |
| 358 | Blaney, C.S.; Mazerolle, D.M. 2010. Fieldwork 2010. Atlantic Canada Conservation Data Centre. Sackville NB, 15508 recs. |
| 324 | Blaney, C.S.; Mazerolle, D.M.; Belliveau, A.B. 2014. Atlantic Canada Conservation Data Centre Fieldwork 2014. Atlantic Canada Conservation Data Centre, # recs. |
| 317 | Scott, F.W. 2002. Nova Scotia Herpetofauna Atlas Database. Acadia University, Wolfville NS, 8856 recs. |
| 306 | Klymko, J. 2018. Maritimes Butterfly Atlas database. Atlantic Canada Conservation Data Centre. |
| 305 | Benjamin, L.K. (compiler). 2012. Significant Habitat & Species Database. Nova Scotia Dept Natural Resources, 4965 recs. |
| 274 | Blaney, C.S.; Mazerolle, D.M. 2012. Fieldwork 2012. Atlantic Canada Conservation Data Centre, 13,278 recs. |
| 265 | Pronych, G. & Wilson, A. 1993. Atlas of Rare Vascular Plants in Nova Scotia. Nova Scotia Museum, Halifax NS, I:1-168, II:169-331. 1446 recs. |
| 263 | Belliveau, A.G. 2018. Atlantic Canada Conservation Data Centre Fieldwork 2017. Atlantic Canada Conservation Data Centre. |
| 252 | Blaney, C.S. & Spicer, C.D.; Popma, T.M.; Basquill, S.P. 2003. Vascular Plant Surveys of Northumberland Strait Rivers & Amherst Area Peatlands. Nova Scotia Museum Research Grant, 501 recs. |
| 247 | Neily, T.H. & Pepper, C.; Toms, B. 2013. Nova Scotia lichen location database. Mersey Tobeatic Research Institute, 1301 records. |
| 244 | Churchill, J.L. 2018. Atlantic Canada Conservation Data Centre Fieldwork 2018. Atlantic Canada Conservation Data Centre, 907 recs. |
| 235 | Chapman-Lam, C.J. 2022. Atlantic Canada Conservation Data Centre 2021 botanical fieldwork. Atlantic Canada Conservation Data Centre, 15099 recs. |
| 211 | Hicks, Andrew. 2009. Coastal Waterfowl Surveys Database, 2000-08. Canadian Wildlife Service, Sackville, 46488 recs (11149 non-zero). |
| 206 | LaPaix, R.W.; Crowell, M.J.; MacDonald, M. 2011. Stantec rare plant records, 2010-11. Stantec Consulting, 334 recs. |
| 198 | Churchill, J.L.; Klymko, J.D. 2015. Chignecto and Tintamarre National Wildlife Area Bird Surveys 2015. Atlantic Canada Conservation Data Centre, 2238 recs. |
| 192 | Toms, B. 2018. Bat Species data from www.batconservation.ca for Nova Scotia. Mersey Tobeatic Research Institute, 547 Records. |
| 180 | Blaney, C.S. & Mazerolle, D.M. 2011. Field data from NCC properties at Musquash Harbour NB & Goose Lake NS. Atlantic Canada Conservation Data Centre, 1739 recs. |
| 178 | Bryson, I. 2013. Nova Scotia rare plant records. CBCL Ltd., 180 records. |
| 149 | Brunelle, P.-M. (compiler). 2009. ADIP/MDDS Odonata Database: data to 2006 inclusive. Atlantic Dragonfly Inventory Program (ADIP), 24200 recs. |
| 140 | Pepper, C. 2013. 2013 rare bird and plant observations in Nova Scotia. , 181 records. |
| 139 | e-Butterfly. 2016. Export of Maritimes records and photos. Maxim Larrivee, Sambo Zhang (ed.) e-butterfly.org. |
| 139 | Neily, T.H. 2017. Nova Scotia lichen records. Mersey Tobeatic Research Institute. |
| 139 | Phinney, Lori. 2020. Pre- and post White-nose Syndrome bat acoustic monitoring, NS. Mersey Tobeatic Research Institute, 1279 recs. |
| 134 | Bryson, I.C. 2020. Nova Scotia flora and lichen observations 2020. Nova Scotia Environment, 139 recs. |
| 129 | Manthorne, A. 2014. MaritimesSwiftwatch Project database 2013-2014. Bird Studies Canada, Sackville NB, 326 recs. |
| 128 | Munro, Marian K. Nova Scotia Provincial Museum of Natural History Herbarium Database. Nova Scotia Provincial Museum of Natural History, Halifax, Nova Scotia. 2013. |
| 127 | MacDonald, Haley. 2022. Updates to <i>Fraxinus nigra</i> observations on NCC Docherty's Brook property. Nature Conservancy of Canada. |
| 126 | Wilhelm, S.I. et al. 2011. Colonial Waterbird Database. Canadian Wildlife Service, Sackville, 2698 sites, 9718 recs (8192 obs). |
| 121 | Blaney, C.S. 2020. Sean Blaney 2020 field data. Atlantic Canada Conservation Data Centre, 4407 records. |
| 116 | Munro, Marian K. Tracked lichen specimens, Nova Scotia Provincial Museum of Natural History Herbarium. Atlantic Canada Conservation Data Centre. 2019. |
| 115 | Blaney, C.S. 2000. Fieldwork 2000. Atlantic Canada Conservation Data Centre. Sackville NB, 1265 recs. |
| 115 | Richardson, Leif. 2018. Maritimes <i>Bombus</i> records from various sources. Richardson, Leif. |
| 114 | Belliveau, A.G. 2021. E.C. Smith Herbarium and Atlantic Canada Conservation Data Centre Fieldwork 2021. E.C. Smith Herbarium. |
| 109 | e-Butterfly. 2019. Export of Maritimes records and photos. McFarland, K. (ed.) e-butterfly.org. |
| 108 | Cameron, R.P. 2009. Cyanolichen database. Nova Scotia Environment & Labour, 1724 recs. |
| 107 | McAlpine, D.F. 1998. NBM Science Collections databases to 1998. New Brunswick Museum, Saint John NB, 241 recs. |
| 107 | Nussey, Pat & NCC staff. 2019. AEI tracked species records, 2016-2019. Chapman, C.J. (ed.) Atlantic Canada Conservation Data Centre, 333. |
| 104 | Chapman, C.J. 2018. Atlantic Canada Conservation Data Centre botanical fieldwork 2018. Atlantic Canada Conservation Data Centre, 11171 recs. |
| 101 | Churchill, J.L.; Walker, J. 2017. Species at Risk Surveys at Correctional Services Canada Properties in Nova Scotia and New Brunswick. Atlantic Canada Conservation Data Centre. |
| 99 | Brazner, J. 2016. Nova Scotia Forested Wetland Bird Surveys. Nova Scotia Department of Lands and Forestry. |
| 98 | iNaturalist. 2018. iNaturalist Data Export 2018. iNaturalist.org and iNaturalist.ca, Web site: 11700 recs. |
| 96 | LaPaix, R.W.; Crowell, M.J.; MacDonald, M.; Neily, T.D.; Quinn, G. 2017. Stantec Nova Scotia rare plant records, 2012-2016. Stantec Consulting. |
| 95 | Belliveau, A.G. 2016. Atlantic Canada Conservation Data Centre Fieldwork 2016. Atlantic Canada Conservation Data Centre, 10695 recs. |
| 93 | Island Nature Trust. 2016. Farmland birds project. Mader, Shannon (ed.) . |
| 88 | Stewart, J.I. 2010. Peregrine Falcon Surveys in New Brunswick, 2002-09. Canadian Wildlife Service, Sackville, 58 recs. |
| 87 | McNeil, J.A. 2018. Wood Turtle records, 2018. Mersey Tobeatic Research Institute, 68 recs. |
| 86 | Chapman-Lam, C.J. 2021. Atlantic Canada Conservation Data Centre 2020 botanical fieldwork. Atlantic Canada Conservation Data Centre, 17309 recs. |
| 86 | Layberry, R.A. & Hall, P.W., LaFontaine, J.D. 1998. The Butterflies of Canada. University of Toronto Press. 280 pp+plates. |
| 85 | Zinck, M. & Roland, A.E. 1998. Roland's Flora of Nova Scotia. Nova Scotia Museum, 3rd ed., rev. M. Zinck; 2 Vol., 1297 pp. |

| # recs | CITATION |
|--------|---|
| 81 | Nature Conservancy of Canada. 2022. NCC Field data for Nova Scotia. Nature Conservancy of Canada. |
| 80 | Staicer, C. 2021. Additional compiled Nova Scotia Species at Risk bird records, 2005-2020. Dalhousie University. |
| 77 | Belland, R.J. Maritimes moss records from various herbarium databases. 2014. |
| 77 | Hagerman, Christianne. 2022. Wisqoq and Eastern White Cedar field work. E.C. Smith Herbarium, Acadia University. |
| 76 | NatureServe Canada. 2019. iNaturalist Maritimes Butterfly Records. iNaturalist.org and iNaturalist.ca. |
| 76 | Tims, J. & Craig, N. 1995. Environmentally Significant Areas in New Brunswick (NBESA). NB Dept of Environment & Nature Trust of New Brunswick Inc, 6042 recs. https://doi.org/10.1037/arc0000014 . |
| 73 | Roland, A.E. & Smith, E.C. 1969. The Flora of Nova Scotia, 1st Ed. Nova Scotia Museum, Halifax, 743pp. |
| 72 | Cameron, R.P. 2011. Lichen observations, 2011. Nova Scotia Environment & Labour, 731 recs. |
| 70 | Blaney, C.S. 2016. Atlantic Canada Conservation Data Centre Fieldwork 2016. Atlantic Canada Conservation Data Centre, 6719 recs. |
| 70 | Bryson, I. 2020. Nova Scotia and Newfoundland rare species observations, 2018-2020. Nova Scotia Environment. |
| 70 | Staicer, C. & Bliss, S.; Achenbach, L. 2017. Occurrences of tracked breeding birds in forested wetlands. , 303 records. |
| 68 | Blaney, C.S.; Mazerolle, D.M.; Belliveau, A.B. 2013. Atlantic Canada Conservation Data Centre Fieldwork 2013. Atlantic Canada Conservation Data Centre, 9000+ recs. |
| 67 | Staicer, Cindy. 2022. 2021 Landbird Species at Risk observations. Dalhousie University. |
| 66 | Blaney, C.S.; Mazerolle, D.M.; Oberndorfer, E. 2007. Fieldwork 2007. Atlantic Canada Conservation Data Centre. Sackville NB, 13770 recs. |
| 65 | Belliveau, A.G. 2014. Plant Records from Southern and Central Nova Scotia. Atlantic Canada Conservation Data Centre, 919 recs. |
| 63 | iNaturalist. 2020. iNaturalist butterfly records selected for the Maritimes Butterfly Atlas. iNaturalist. |
| 62 | Blaney, C.S.; Mazerolle, D.M.; Hill, N.M. 2011. Nova Scotia Crown Share Land Legacy Trust Fieldwork. Atlantic Canada Conservation Data Centre, 5022 recs. |
| 62 | Toms, Brad & Pepper, Chris; Neily, Tom. 2022. Nova Scotia lichen database [as of 2022-04]. Mersey Tobeatic Research Institute. |
| 60 | Belliveau, A.G. 2018. E.C. Smith Herbarium and Atlantic Canada Conservation Data Centre Fieldwork 2018. E.C. Smith Herbarium, 6226 recs. |
| 52 | Blaney, C.S. 2018. Atlantic Canada Conservation Data Centre Fieldwork 2018. Atlantic Canada Conservation Data Centre. |
| 51 | Blaney, C.S.; Spicer, C.D.; Rothfels, C. 2004. Fieldwork 2004. Atlantic Canada Conservation Data Centre. Sackville NB, 1343 recs. |
| 51 | Churchill, J.L. 2020. Atlantic Canada Conservation Data Centre Fieldwork 2020. Atlantic Canada Conservation Data Centre, 1083 recs. |
| 50 | Hall, R.A. 2003. NS Freshwater Mussel Fieldwork. Nova Scotia Dept Natural Resources, 189 recs. |
| 49 | Hall, R.A. 2001. S.. NS Freshwater Mussel Fieldwork. Nova Scotia Dept Natural Resources, 178 recs. |
| 48 | Amirault, D.L. & McKnight, J. 2003. Piping Plover Database 1991-2003. Canadian Wildlife Service, Sackville, unpublished data. 7 recs. |
| 48 | Porter, Caitlin et al. 2022. 2021 wildlife observations from the Caemmerer NCC site. Atlantic Canada Conservation Data Centre. |
| 47 | Blaney, C.S.; Spicer, C.D.; Popma, T.M.; Hanel, C. 2002. Fieldwork 2002. Atlantic Canada Conservation Data Centre. Sackville NB, 2252 recs. |
| 47 | Patrick, Allison. 2021. Animal and plant records from NCC properties from 2019 and 2020. Nature Conservancy Canada. |
| 46 | Chapman, C.J. 2019. Atlantic Canada Conservation Data Centre 2019 botanical fieldwork. Atlantic Canada Conservation Data Centre, 11729 recs. |
| 46 | Klymko, J.J.D. 2012. Insect fieldwork & submissions, 2011. Atlantic Canada Conservation Data Centre. Sackville NB, 760 recs. |
| 46 | Spicer, C.D. & Harries, H. 2001. Mount Allison Herbarium Specimens. Mount Allison University, 128 recs. |
| 45 | Blaney, C.S. 2003. Fieldwork 2003. Atlantic Canada Conservation Data Centre. Sackville NB, 1042 recs. |
| 45 | Cameron, R.P. 2009. Erioderma pedicellatum database, 1979-2008. Dept Environment & Labour, 103 recs. |
| 45 | Canadian Wildlife Service, Dartmouth. 2010. Piping Plover censuses 2007-09, 304 recs. |
| 45 | Hubley, Nicole. 2022. Monarch (<i>Danaus plexippus</i>) records submitted to MTRI from the 2021 field season. Mersey Tobeatic Research Institute. |
| 43 | Blaney, C.S.; Spicer, C.D. 2001. Fieldwork 2001. Atlantic Canada Conservation Data Centre. Sackville NB, 981 recs. |
| 42 | Benjamin, L.K. (compiler). 2001. Significant Habitat & Species Database. Nova Scotia Dept of Natural Resources, 15 spp, 224 recs. |
| 40 | Cameron, E. 2007. Canadian Gypsum Co. survey 2005-07. Dillon Consulting Ltd, 40 recs. |
| 40 | Neily, T.H. & Pepper, C.; Toms, B. 2020. Nova Scotia lichen database [as of 2020-03-18]. Mersey Tobeatic Research Institute. |
| 39 | MacDonald, E.C. 2018. CWS Piping Plover Census, 2010-2017. Canadian Wildlife Service, 672 recs. |
| 39 | Mazerolle, D.M. 2018. Atlantic Canada Conservation Data Centre botanical fieldwork 2018. Atlantic Canada Conservation Data Centre, 13515 recs. |
| 38 | Blaney, C.S. 2017. Atlantic Canada Conservation Data Centre Fieldwork 2017. Atlantic Canada Conservation Data Centre. |
| 38 | Nova Scotia Nature Trust. 2013. Nova Scotia Nature Trust 2013 Species records. Nova Scotia Nature Trust, 95 recs. |
| 38 | Thomas, P. 2018. CSC Dorchester Bobolink Survey. Environment Canada, Canadian Wildlife Service. |
| 37 | Blaney, C.S.; Mazerolle, D.M.; Klymko, J.; Spicer, C.D. 2006. Fieldwork 2006. Atlantic Canada Conservation Data Centre. Sackville NB, 8399 recs. |
| 37 | Klymko, J. 2016. Atlantic Canada Conservation Data Centre Fieldwork 2016. Atlantic Canada Conservation Data Centre. |
| 37 | Neily, T.H. & Pepper, C.; Toms, B. 2015. Nova Scotia lichen location database [as of 2015-02-15]. Mersey Tobeatic Research Institute, 1691 records. |
| 36 | Blaney, C.S.; Mazerolle, D.M. 2008. Fieldwork 2008. Atlantic Canada Conservation Data Centre. Sackville NB, 13343 recs. |
| 36 | Churchill, J.L. 2018. Atlantic Canada Conservation Data Centre Fieldwork 2017. Atlantic Canada Conservation Data Centre, 2318 recs. |
| 36 | Klymko, J.J.D.; Robinson, S.L. 2012. 2012 field data. Atlantic Canada Conservation Data Centre, 447 recs. |
| 36 | Mazerolle, D.M. 2017. Atlantic Canada Conservation Data Centre Fieldwork 2017. Atlantic Canada Conservation Data Centre. |
| 35 | Mersey Tobeatic Research Institute. 2021. 2020 Monarch records from the MTRI monitoring program. Mersey Tobeatic Research Institute, 72 records. |
| 35 | Ogden, J. NS DNR Butterfly Collection Dataset. Nova Scotia Department of Natural Resources. 2014. |
| 35 | Pepper, C. 2021. Rare bird, plant and mammal observations in Nova Scotia, 2017-2021. |
| 35 | Porter, C.J.M. 2014. Field work data 2007-2014. Nova Scotia Nature Trust, 96 recs. |
| 33 | Arsenault, M. 2019. Cormorant colony nest counts. PE Department of Communities, Land, and Environment. |
| 32 | Glen, W. 1991. 1991 Prince Edward Island Forest Biomass Inventory Data. PEI Dept of Energy and Forestry, 10059 recs. |
| 31 | Neily, T.H. 2019. Tom Neily NS Bryophyte records (2009-2013). T.H. Neily, Atlantic Canada Conservation Data Centre, 1029 specimen records. |
| 30 | Erskine, A.J. 1999. Maritime Nest Records Scheme (MNRS) 1937-1999. Canadian Wildlife Service, Sackville, 313 recs. |

| # recs | CITATION |
|--------|---|
| 30 | Ferguson, D.C. 1954. The Lepidoptera of Nova Scotia. Part I, macrolepidoptera. Proceedings of the Nova Scotian Institute of Science, 23(3), 161-375. |
| 29 | Wallace, S. 2020. Stewardship Department species occurrence data on NTNB preserves. Nature Trust of New Brunswick. |
| 29 | Westwood, A., Staicer, C. 2016. Nova Scotia landbird Species at Risk observations. Dalhousie University. |
| 28 | Pepper, Chris. 2012. Observations of breeding Canada Warbler's along the Eastern Shore, NS. Pers. comm. to S. Blaney, Jan. 20, 28 recs. |
| 26 | Cameron, R.P. 2018. <i>Degelia plumbea</i> records. Nova Scotia Environment. |
| 26 | Porter, Caitlin. 2021. Field data for 2020 in various locations across the Maritimes. Atlantic Canada Conservation Data Centre, 3977 records. |
| 25 | Haughian, Sean. 2021. Update to lichen data from 2017-2021. Nova Scotia Museum. |
| 25 | MacDonald, E.C. 2018. Piping Plover nest records from 2010-2017. Canadian Wildlife Service. |
| 25 | Popma, T.M. 2003. Fieldwork 2003. Atlantic Canada Conservation Data Centre. Sackville NB, 113 recs. |
| 24 | Belliveau, A.G. 2021. New Black ash site records near Kentville, NS. Acadia University, 47 records. |
| 24 | LaPaix, Rich. 2022. Rare species observations, 2018-2022. Nova Scotia Nature Trust. |
| 24 | Powell, B.C. 1967. Female sexual cycles of <i>Chrysemy spicta</i> & <i>Clemmys insculpta</i> in Nova Scotia. Can. Field-Nat., 81:134-139. 26 recs. |
| 23 | Benjamin, L.K. 2011. NSDNR fieldwork & consultant reports 1997, 2009-10. Nova Scotia Dept Natural Resources, 85 recs. |
| 23 | Catling, P.M., Erskine, D.S. & MacLaren, R.B. 1985. The Plants of Prince Edward Island with new records, nomenclatural changes & corrections & deletions, 1st Ed. Research Branch, Agriculture Canada, Ottawa, Publication 1798. 22pp. |
| 23 | Manthorne, A. 2019. Incidental aerial insectivore observations. Birds Canada. |
| 23 | Ogden, K. Nova Scotia Museum butterfly specimen database. Nova Scotia Museum. 2017. |
| 22 | Klymko, John. 2022. Atlantic Canada Conservation Data Centre zoological fieldwork 2021. Atlantic Canada Conservation Data Centre. |
| 22 | Nelly, T.H. 2006. <i>Cypripedium arietinum</i> in Hants Co. Pers. comm. to C.S. Blaney. 22 recs, 22 recs. |
| 22 | NS DNR. 2017. Black Ash records from NS DNR Permanent Sample Plots (PSPs), 1965-2016. NS Dept of Natural Resources. |
| 21 | Archibald, D.R. 2003. NS Freshwater Mussel Fieldwork. Nova Scotia Dept Natural Resources, 213 recs. |
| 21 | Belliveau, A. 2013. Rare species records from Nova Scotia. Mersey Tobeatic Research Institute, 296 records. 296 recs. |
| 21 | Churchill, J.L., Klymko, J.D.D. 2016. Atlantic Canada Conservation Data Centre Fieldwork 2016. Atlantic Canada Conservation Data Centre. |
| 20 | Blaney, C.S.; Spicer, C.D.; Mazerolle, D.M. 2005. Fieldwork 2005. Atlantic Canada Conservation Data Centre. Sackville NB, 2333 recs. |
| 20 | Feltham, Carter. 2022. Monarch (<i>Danaus plexippus</i>) and Milkweed MTRI records from the 2022 Field Season. Mersey Tobeatic Research Institute. |
| 20 | Klymko, J.J.D. 2016. 2015 field data. Atlantic Canada Conservation Data Centre. |
| 20 | Neily, T.H. 2013. Email communication to Sean Blaney regarding <i>Listera australis</i> observations made from 2007 to 2011 in Nova Scotia. , 50. |
| 20 | Phinney, Lori; Toms, Brad; et. al. 2016. Bank Swallows (<i>Riparia riparia</i>) in Nova Scotia: inventory and assessment of colonies. Merser Tobeatic Research Institute, 25 recs. |
| 19 | Godbout, V. 2002. SAR Inventory: Birds in Fort Beauséjour NHS. Parks Canada, Atlantic, SARINV02-01. 202 recs. |
| 18 | Burns, L. 2013. Personal communication concerning bat occurrence on PEI. Winter 2013. Pers. comm. |
| 18 | Cameron, R.P. 2014. 2013-14 rare species field data. Nova Scotia Department of Environment, 35 recs. |
| 18 | Klymko, J.J.D. 2018. 2017 field data. Atlantic Canada Conservation Data Centre. |
| 18 | Munro, Marian K. Nova Scotia Provincial Museum of Natural History Herbarium Database. Nova Scotia Provincial Museum of Natural History, Halifax, Nova Scotia. 2014. |
| 17 | Anderson, Frances; Neily, Tom. 2010. A Reconnaissance Level Survey of Calciphilous Lichens in Selected Karst Topography in Nova Scotia with Notes on Incidental Bryophytes. Mersey Tobeatic Research Institute. |
| 17 | McNeil, J.A. 2016. Blandings Turtle (<i>Emydoidea blandingii</i>), Eastern Ribbonsnake (<i>Thamnophis sauritus</i>), Wood Turtle (<i>Glyptemys insculpta</i>), and Snapping Turtle (<i>Chelydra serpentina</i>) sightings, 2016. Mersey Tobeatic Research Institute, 774 records. |
| 17 | Neily, T.H. 2010. <i>Erioderma pedicellatum</i> records 2005-09. Mersey Tobiatic Research Institute, 67 recs. |
| 17 | Richardson, D., Anderson, F., Cameron, R, McMullin, T., Clayden, S. 2014. Field Work Report on Black Foam Lichen (<i>Anzia colpodes</i>). COSEWIC. |
| 16 | Blaney, C.S.; Mazerolle, D.M. 2009. Fieldwork 2009. Atlantic Canada Conservation Data Centre. Sackville NB, 13395 recs. |
| 16 | Neily, T.H. & Pepper, C. 2020. Nova Scotia SMP lichen surveys 2020. Mersey Tobeatic Research Institute. |
| 15 | Basquill, S.P. 2011 vascular plant field data. Nova Scotia Department of Natural Resources, 37 recs. |
| 15 | Blaney, C.S. 2019. Sean Blaney 2019 field data. Atlantic Canada Conservation Data Centre, 4407 records. |
| 15 | Cameron, R.P. 2013. 2013 rare species field data. Nova Scotia Department of Environment, 71 recs. |
| 15 | Chiasson, R. 2018. Breeding bird observations from NBWTF project. pers. comm. to S. Blaney. |
| 14 | Benedict, B. Connell Herbarium Specimens (Data) . University New Brunswick, Fredericton. 2003. |
| 14 | Blaney, C.S.; Mazerolle, D.M. 2011. Fieldwork 2011. Atlantic Canada Conservation Data Centre. Sackville NB. |
| 13 | e-Butterfly. 2018. Selected Maritimes butterfly records from 2016 and 2017. Maxim Larrivee, Sambo Zhang (ed.) e-butterfly.org. |
| 13 | Gilhen, J. 1984. Amphibians & Reptiles of Nova Scotia, 1st Ed. Nova Scotia Museum, 164pp. |
| 13 | McMullin, R.T. 2022. Maritimes lichen records. Canadian Museum of Nature. |
| 13 | Neily, T.H. 2012. 2012 <i>Erioderma pedicellatum</i> records in Nova Scotia. |
| 13 | Nova Scotia Nature Trust. 2014. Ladyslipper records from Saint Croix Nova Scotia, JLC Ed. Nova Scotia Nature Trust. |
| 12 | Basquill, S.P. 2012. 2012 rare vascular plant field data. Nova Scotia Department of Natural Resources, 37 recs. |
| 12 | Cameron, R.P. 2017. 2017 rare species field data. Nova Scotia Environment, 64 recs. |
| 12 | Chaput, G. 2002. Atlantic Salmon: Maritime Provinces Overview for 2001. Dept of Fisheries & Oceans, Atlantic Region, Science Stock Status Report D3-14. 39 recs. |
| 12 | Clayden, S.R. 1998. NBM Science Collections databases: vascular plants. New Brunswick Museum, Saint John NB, 19759 recs. |
| 12 | Klymko, J. Dataset of butterfly records at the New Brunswick Museum not yet accessioned by the museum. Atlantic Canada Conservation Data Centre. 2016. |
| 12 | Oldham, M.J. 2000. Oldham database records from Maritime provinces. Oldham, M.J; ONHIC, 487 recs. |
| 11 | Bateman, M.C. 2001. Coastal Waterfowl Surveys Database, 1965-2001. Canadian Wildlife Service, Sackville, 667 recs. |
| 11 | Doucet, D.A. 2009. Census of Globally Rare, Endemic Butterflies of Nova Scotia Gulf of St Lawrence Salt Marshes. Nova Scotia Dept of Natural Resources, Species at Risk, 155 recs. |

| # recs | CITATION |
|--------|---|
| 11 | Hill, N.M. 1994. Status report on the Long's bulrush <i>Scirpus longii</i> in Canada. Committee on the Status of Endangered Wildlife in Canada, 7 recs. |
| 11 | Robinson, S.L. 2015. 2014 field data. |
| 10 | Adams, J. & Herman, T.B. 1998. Thesis, Unpublished map of <i>C. insculpta</i> sightings. Acadia University, Wolfville NS, 88 recs. |
| 10 | Basquill, S.P. 2003. Fieldwork 2003. Atlantic Canada Conservation Data Centre, Sackville NB, 69 recs. |
| 10 | Belliveau, A.G. & Vail, Cole; King, Katie. 2020. New <i>Allium tricoccum</i> locations, Cornwallis River. Chapman, C.J. (ed.) Acadia University. |
| 10 | Benedict, B. Connell Herbarium Specimens. University New Brunswick, Fredericton. 2003. |
| 10 | Doucet, D.A. 2007. Lepidopteran Records, 1988-2006. Doucet, 700 recs. |
| 10 | Downes, C. 1998-2000. Breeding Bird Survey Data. Canadian Wildlife Service, Ottawa, 111 recs. |
| 10 | Goltz, J.P. & Bishop, G. 2005. Confidential supplement to Status Report on Prototype Quillwort (<i>Isoetes prototypus</i>). Committee on the Status of Endangered Wildlife in Canada, 111 recs. |
| 10 | Neily, T. H. 2018. Lichen and Bryophyte records, AEI 2017-2018. Tom Neily; Atlantic Canada Conservation Data Centre. |
| 9 | Benjamin, L.K. 2006. <i>Cyrtopodium arietinum</i> . Pers. comm. to D. Mazerolle. 9 recs, 9 recs. |
| 9 | Benjamin, L.K. 2012. NSDNR fieldwork & consultant reports 2008-2012. Nova Scotia Dept Natural Resources, 196 recs. |
| 9 | Cameron, R.P. 2005. <i>Erioderma pedicellatum</i> unpublished data. NS Dept of Environment, 9 recs. |
| 9 | Cameron, R.P. 2006. <i>Erioderma pedicellatum</i> 2006 field data. NS Dept of Environment, 9 recs. |
| 9 | Chapman, C.N. (Cody). 2020. Nova Scotia Black Ash (<i>Fraxinus nigra</i>) field observations by Confederacy of Mainland Mi'kmaq. Forestry Program, Confederacy of Mainland Mi'kmaq. |
| 9 | Clayden, S.R. 2007. NBM Science Collections databases: vascular plants. New Brunswick Museum, Saint John NB, download Mar. 2007, 6914 recs. |
| 9 | Epworth, W. 2012. Species at Risk records, 2009-11. Fort Folly Habitat Recovery Program, 162 recs. |
| 9 | Hughes, Cory. 2020. Atlantic Forestry Centre <i>Coccinella transversoguttata</i> collections. Canadian Forest Service, Atlantic Forestry Centre. |
| 9 | Neily, T.H. & Pepper, C.; Toms, B. 2020. Nova Scotia lichen database [as of 2020-05-25]. Mersey Tobeatic Research Institute, 668 recs. |
| 9 | Sollows, M.C., 2009. NBM Science Collections databases: molluscs. New Brunswick Museum, Saint John NB, download Jan. 2009, 6951 recs (2957 in Atlantic Canada). |
| 9 | Webster, R.P. Atlantic Forestry Centre Insect Collection, Maritimes butterfly records. Natural Resources Canada. 2014. |
| 8 | Holder, M.L.; Kingsley, A.L. 2000. Kinglsey and Holder observations from 2000 field work. |
| 8 | King, Katie; Jean, Samuel. 2021. Black ash observations near Booklyn, NS. E.C. Smith Herbarium. |
| 8 | Sollows, M.C., 2008. NBM Science Collections databases: mammals. New Brunswick Museum, Saint John NB, download Jan. 2008, 4983 recs. |
| 7 | Blaney, C.S. Miscellaneous specimens received by ACCDC (botany). Various persons. 2001-08. |
| 7 | Cameron, B. 2006. <i>Hepatica americana</i> Survey at Scotia Mine Site in Gays River, and Discovery of Three Yellow-listed Species. Conestoga-Rovers and Associates, (a consulting firm), october 25. 7 recs. |
| 7 | O'Neil, S. 1998. Atlantic Salmon: Northumberland Strait Nova Scotia part of SFA 18. Dept of Fisheries & Oceans, Atlantic Region, Science. Stock Status Report D3-08. 9 recs. |
| 7 | Olsen, R. Herbarium Specimens. Nova Scotia Agricultural College, Truro. 2003. |
| 7 | Speers, L. 2008. Butterflies of Canada database: New Brunswick 1897-1999. Agriculture & Agri-Food Canada, Biological Resources Program, Ottawa, 2048 recs. |
| 6 | Benjamin, L.K. 2009. Boreal Felt Lichen, Mountain Avens, Orchid and other recent records. Nova Scotia Dept Natural Resources, 105 recs. |
| 6 | Boyne, A.W. & Grecian, V.D. 1999. Tern Surveys. Canadian Wildlife Service, Sackville, unpublished data. 23 recs. |
| 6 | Brazner, J.; Hill, N. 2018. Plant observations along the Cornwallis River, Nova Scotia. Nova Scotia Department of Lands and Forestry. |
| 6 | eBird. 2021. eBird Basic Dataset. Version: EBD_relOct-2020. Ithaca, New York. Oct 2020, Prince Edward Island Bird SAR subset. Cornell Lab of Ornithology. |
| 6 | Edsall, J. 2007. Personal Butterfly Collection: specimens collected in the Canadian Maritimes, 1961-2007. J. Edsall, unpubl. report, 137 recs. |
| 6 | Hall, R. 2008. Rare plant records in old fieldbook notes from Truro area. Pers. comm. to C.S. Blaney. 6 recs, 6 recs. |
| 6 | Klymko, J. 2019. Atlantic Canada Conservation Data Centre zoological fieldwork 2018. Atlantic Canada Conservation Data Centre. |
| 6 | Klymko, J. Butterfly records at the Nova Scotia Museum not yet accessioned by the museum. Atlantic Canada Conservation Data Centre. 2017. |
| 6 | Matthew Smith. 2010. Field trip report from Avon Caving Club outlining the discovery of <i>Cyrtopodium arietinum</i> and <i>Hepatica nobilis</i> populations. Public Works and Government Services Canada. |
| 6 | Mazerolle, D.M. 2020. Atlantic Canada Conservation Data Centre botanical fieldwork 2019. Atlantic Canada Conservation Data Centre. |
| 6 | Neily, T.H. Tom Neily NS Sphagnum records (2009-2014). T.H. Neily, Atlantic Canada Conservation Data Centre. 2019. |
| 6 | Nova Scotia Nature Trust. 2022. Ram's Head Lady Slipper observations from 2015 and 2019. , 6 records. |
| 6 | Pike, E., Tingley, S. & Christie, D.S. 2000. Nature NB Listserve. University of New Brunswick, listserv.unb.ca/archives/naturenb. 68 recs. |
| 6 | Richardson, D., Anderson, F., Cameron, R, Pepper, C., Clayden, S. 2015. Field Work Report on the Wrinkled Shingle lichen (<i>Pannaria lurida</i>). COSEWIC. |
| 6 | White, S. 2019. Notable species sightings, 2018. East Coast Aquatics. |
| 5 | Bagnell, B.A. 2001. New Brunswick Bryophyte Occurrences. B&B Botanical, Sussex, 478 recs. |
| 5 | Cameron, R.P. 2012. Additional rare plant records, 2009. , 7 recs. |
| 5 | Carter, Jeff; Churchill, J.; Churchill, I.; Churchill, L. 2020. Bank Swallow colony Scots Bay, NS. Atlantic Canada Conservation Data Centre. |
| 5 | Erskine, D. 1960. The plants of Prince Edward Island, 1st Ed. Research Branch, Agriculture Canada, Ottawa., Publication 1088. 1238 recs. |
| 5 | Gallop, John. 2021. Sheet Harbour rare lichen observations. McCallum Environmental. |
| 5 | Klymko, J. 2021. Atlantic Canada Conservation Data Centre zoological fieldwork 2020. Atlantic Canada Conservation Data Centre. |
| 5 | Mazerolle, D.M. 2016. Atlantic Canada Conservation Data Centre Fieldwork 2017. Atlantic Canada Conservation Data Centre. |
| 5 | McNeil, J.A. 2020. Snapping Turtle and Eastern Painted Turtle records, 2020. Mersey Tobeatic Research Institute. |
| 5 | Sabine, D.L. 2013. Dwaine Sabine butterfly records, 2009 and earlier. |
| 5 | Tingley, S. (compiler). 2001. Butterflies of New Brunswick. , Web site: www.geocities.com/Yosemite/8425/buttrfly. 142 recs. |
| 5 | Towell, C. 2014. 2014 Northern Goshawk and Common Nighthawk email reports, NS. NS Department of Natural Resources. |
| 5 | Walker, J. 2017. Bird inventories at French River, NS, and Memramcook, NB, for Nature Conservancy of Canada. Pers. comm. to AC CDC. |
| 5 | Webster, R.P. & Edsall, J. 2007. 2005 New Brunswick Rare Butterfly Survey. Environmental Trust Fund, unpublished report, 232 recs. |
| 4 | Basquill, S.P., Porter, C. 2019. Bryophyte and lichen specimens submitted to the E.C. Smith Herbarium. NS Department of Lands and Forestry. |
| 4 | Benjamin, L.K. 2009. NSDNR Fieldwork & Consultants Reports. Nova Scotia Dept Natural Resources, 143 recs. |

| # recs | CITATION |
|--------|---|
| 4 | Bredin, K.A. 2002. NS Freshwater Mussel Fieldwork. Atlantic Canada Conservation Data Centre, 30 recs. |
| 4 | Cameron-MacMillan, Maureen. 2020. Northern Goshawk Nests in Eastern Nova Scotia, as of November, 2020. Nova Scotia Department of Lands and Forestry. |
| 4 | Clayden, S.R. 2005. Confidential supplement to Status Report on Ghost Antler Lichen (<i>Pseudevernia cladonia</i>). Committee on the Status of Endangered Wildlife in Canada, 27 recs. |
| 4 | Cody, W.J. 2003. Nova Scotia specimens of <i>Equisetum pratense</i> at the DAO herbarium in Ottawa. , Pers. comm. to C.S. Blaney. 4 recs. |
| 4 | Curley, F.R. 2005. PEF&W Collection 2003-04. PEI Fish & Wildlife Div., 716 recs. |
| 4 | Forsythe, B. 2006. <i>Cypripedium arietinum</i> at Meadow Pond, Hants Co. Pers. comm. to C.S. Blaney. 4 recs, 4 recs. |
| 4 | Klymko, J.J.D.; Robinson, S.L. 2014. 2013 field data. Atlantic Canada Conservation Data Centre. |
| 4 | Mills, Pamela. 2007. <i>Iva frutescens</i> records. Nova Scotia Dept of Natural Resources, Wildlife Div. Pers. comm. to S. Basquill, 4 recs. |
| 4 | Plissner, J.H. & Haig, S.M. 1997. 1996 International piping plover census. US Geological Survey, Corvallis OR, 231 pp. |
| 3 | Basquill, S.P. 2009. 2009 field observations. Nova Scotia Dept of Natural Resources. |
| 3 | Basset, I.J. & Crompton, C.W. 1978. The Genus <i>Suaeda</i> (<i>Chenopodiaceae</i>) in Canada. Canadian Journal of Botany, 56: 581-591. |
| 3 | Belland, R.J. 2012. PEI moss records from Devonian Botanical Garden. DBG Cryptogam Database, Web site: https://secure.devonian.ualberta.ca/bryo_search.php 748 recs. |
| 3 | Brunelle, P.-M. (compiler). 2010. ADIP/MDDS Odonata Database: NB, NS Update 1900-09. Atlantic Dragonfly Inventory Program (ADIP), 935 recs. |
| 3 | Calhoun, J.C. Butterfly records databased at the McGuire Center for Lepidoptera and Biodiversity. Calhoun, J.C. 2020. |
| 3 | Cameron, R.P. 2012. Rob Cameron 2012 vascular plant data. NS Department of Environment, 30 recs. |
| 3 | Chapman, Cody. Unreported Species at Risk Records across Nova Scotia. Chapman, Cody, 5 records. |
| 3 | Chiasson, Roland. 2017. New Brunswick Breeding bird observations from NBWTF project. pers. comm. to J. Churchill. |
| 3 | Christie, D.S. 2000. Christmas Bird Count Data, 1997-2000. Nature NB, 54 recs. |
| 3 | Curley, F.R. 2007. PEF&W Collection. PEI Fish & Wildlife Div., 199 recs. |
| 3 | Daury, R.W. & Bateman, M.C. 1996. The Barrow's Goldeneye (<i>Bucephala islandica</i>) in the Atlantic Provinces and Maine. Canadian Wildlife Service, Sackville, 47pp. |
| 3 | Doubt, J. 2013. Email to Sean Blaney with Nova Scotia records of <i>Fissidens exilis</i> at Canadian Museum of Nature. pers. comm., 3 records. |
| 3 | Edsall, J. 2001. Lepidopteran records in New Brunswick, 1997-99. , Pers. comm. to K.A. Bredin. 91 recs. |
| 3 | Frittaion, C. 2012. NSNT 2012 Field Observations. Nova Scotia Nature Trust, Pers comm. to S. Blaney Feb. 7, 34 recs. |
| 3 | Hill, N. and D. Patriquin. 2013. 2013 rare plant observations in Williams Lake Backlands area. Fern Hill Institute of Plant Conservation, Berwick, Nova Scotia, 3 records. |
| 3 | Hinds, H.R. 1986. Notes on New Brunswick plant collections. Connell Memorial Herbarium, unpubl, 739 recs. |
| 3 | Klymko, J.J.D. 2011. Insect fieldwork & submissions, 2010. Atlantic Canada Conservation Data Centre. Sackville NB, 742 recs. |
| 3 | McNeil, J.A. 2019. Snapping Turtle records, 2019. Mersey Tobeatic Research Institute. |
| 3 | Mersey Tobeatic Research Institute. 2022. Nova Scotia Bobolink observations. pers. comm. to J. Churchill. |
| 3 | Nature Trust of New Brunswick. 2021. Nature Trust of New Brunswick site inventory data submitted in April 2021. Nature Trust of New Brunswick, 2189 records. |
| 3 | O'Neil, S. 1998. Atlantic Salmon: Eastern Shore Nova Scotia SFA 20. Dept of Fisheries & Oceans, Atlantic Region, Science. Stock Status Report D3-10. 4 recs. |
| 3 | Parker, M. 2016. Wood turtle (<i>Glyptemys insculpta</i>) Visual Surveys at Black, Wallace, Musquodobit and Sackville Rivers, Nova Scotia. East Coast Aquatics Inc., 3 records. |
| 3 | Sabine, M. 2016. NB DNR staff incidental Black Ash observations. New Brunswick Department of Natural Resources. |
| 3 | Standley, L.A. 2002. <i>Carex haydenii</i> in Nova Scotia. , Pers. comm. to C.S. Blaney. 4 recs. |
| 3 | Thompson, R. 2018. Williamsdale Quarry Expansion Project, NS, Environmental Assessment rare plants. Dexter Construction Company Limited. |
| 3 | Wilhelm, S.I. et al. 2019. Colonial Waterbird Database. Canadian Wildlife Service. |
| 2 | Amirault, D.L. 1997-2000. Unpublished files. Canadian Wildlife Service, Sackville, 470 recs. |
| 2 | Amirault, D.L. 2003. 2003 Peregrine Falcon Survey. Canadian Wildlife Service, Sackville, unpublished data. 7 recs. |
| 2 | Amiro, Peter G. 1998. Atlantic Salmon: Inner Bay of Fundy SFA 22 & part of SFA 23. Dept of Fisheries & Oceans, Atlantic Region, Science Stock Status Report D3-12. 4 recs. |
| 2 | Atlantic Canada Bank Swallow Working Group. 2022. 2021 Bank Swallow colony records. Birds Canada. |
| 2 | Bredin, K.A. 2001. WTF Project: Freshwater Mussel Fieldwork in Freshwater Species data. Atlantic Canada Conservation Data Centre, 101 recs. |
| 2 | Cameron, B. 2005. <i>C. palmicola</i> , <i>E. pedicellatum</i> records from Sixth Lake. Pers. comm. to C.S. Blaney. 3 recs, 3 recs. |
| 2 | Cameron, R.P. 2009. Nova Scotia nonvascular plant observations, 1995-2007. Nova Scotia Dept Natural Resources, 27 recs. |
| 2 | de Graaf, M.; Miller, D. 2020. Records of <i>Cypripedium reginae</i> and <i>Equisetum variegatum</i> from CFI property at Scoudouc Road and <i>Symplocarpus foetidus</i> from CFI properties at Upper Cape, Westmorland County, NB. pers. comm. (ed.) Community Forests International. 4 records. |
| 2 | Dibblee, R.L. 1999. PEI Cormorant Survey. Prince Edward Island Fisheries, Aquaculture & Environment, 1p. 21 recs. |
| 2 | Golder Associates. 2018. Dorchester wind turbine bat detections. Owens, Luke, Firman, Mitch, Melcher, Heather (ed.) Golder Associates Ltd. |
| 2 | Harding, R.W. 2008. Harding Personal Insect Collection 1999-2007. R.W. Harding, 309 recs. |
| 2 | Hicklin, P.W. 1995. The Maritime Shorebird Survey Newsletter. Calidris, No. 3. 6 recs. |
| 2 | Honeyman, K. 2019. Unique Areas Database, 2018. J.D. Irving Ltd. |
| 2 | iNaturalist.ca. 2022. iNaturalist records 2022. iNaturalist.ca (ed.) iNaturalist.org; iNaturalist.ca. |
| 2 | Kennedy, Joseph. 2010. New Brunswick Peregrine records, 2009. New Brunswick Dept Natural Resources, 19 recs (14 active). |
| 2 | Kennedy, Joseph. 2010. New Brunswick Peregrine records, 2010. New Brunswick Dept Natural Resources, 16 recs (11 active). |
| 2 | Klymko, J.J.D. 2010. Miscellaneous observations reported to ACCDC (zoology). Pers. comm. from various persons, 3 recs. |
| 2 | LaPaix, R.; Parker, M. 2013. email to Sean Blaney regarding <i>Listera australis</i> observations near Kearney Lake. East Coast Aquatics, 2. |
| 2 | Macaulay, M. Notes on newly discovered <i>Hepatica nobilis</i> var. <i>obtusata</i> population in Cumberland Co. NS. Pers. comm. to S. Blaney, 1 rec. |
| 2 | Mazerolle, D.M. 2005. Bouctouche Irving Eco-Centre rare coastal plant fieldwork results 2004-05. Irving Eco-centre, la Dune du Bouctouche, 174 recs. |
| 2 | Mazerolle, David. 2021. Botanical fieldwork 2019-20200. Parks Canada. |
| 2 | McLean, K. 2019. Species At Risk observations. Clean Annapolis River Project. |
| 2 | McLean, K. 2020. Species occurrence records from Clean Annapolis River Project fieldwork in 2020. Clean Annapolis River Project, 206 records. |

| # recs | CITATION |
|--------|---|
| 2 | Munro, M. 2003. <i>Caulophyllum thalictroides</i> & <i>Carex hirtifolia</i> at Herbert River, Brooklyn, NS. , Pers. comm. to C.S. Blaney. 2 recs. |
| 2 | Munro, M. 2003. <i>Dorca palustris</i> & <i>Hepatica nobilis</i> var. <i>obtusa</i> at Cogmagun River, NS. , Pers. comm. to C.S. Blaney . 2 recs. |
| 2 | Neily, T.H.; Smith, C.; Whitman, E. 2011. NCC Logging Lake (Halifax Co. NS) properties baseline survey data. Nature Conservancy of Canada, 2 recs. |
| 2 | Neily, Tom. 2020. Lichen surveys for PEI Forested Landscapes Priority Place. Chapman, C.J. (ed.) Atlantic Canada Conservation Data Centre, 158 records. |
| 2 | Newell, R. E., MacKinnon, C. M. & Kennedy, A. C. 2006. Botanical Survey of Boot Island National Wildlife Area, Nova Scotia, 2004. Canadian Wildlife Service, Atlantic Region, Technical Report Series Number 450. 3 recs. |
| 2 | Newell, R.E. 2006. Rare plant observations in Digby Neck. Pers. comm. to S. Blaney, 6 recs. |
| 2 | Nova Scotia Department of Lands and Forestry. 2018. Wood Turtle observations in, or near, the cornwallis River watershed. NS DLF, pers. comm. to AC CDC. |
| 2 | Phillips, B. 2017. Emails to John Klymko regarding Eastern Waterfan (<i>Peltigera hydrothyria</i>) occurrences in Fundy National Park. Fundy Biosphere Reserve, 3 recs. |
| 2 | Pulsifer, M.D. 2002. NS Freshwater Mussel Fieldwork. Nova Scotia Dept Natural Resources, 369 recs. |
| 2 | Robinson, S.L. 2010. Fieldwork 2009 (dune ecology). Atlantic Canada Conservation Data Centre. Sackville NB, 408 recs. |
| 2 | Robinson, S.L. 2011. 2011 ND dune survey field data. Atlantic Canada Conservation Data Centre, 2715 recs. |
| 2 | Shafer, A.B.A., D.T. Stewart. 2006. A Disjunct Population of <i>Sorex dispar</i> (Long-Tailed Shrew) in Nova Scotia. <i>Northeastern Naturalist</i> , 13(4): 603-608. |
| 2 | Shortt, R. UNB specimen data for various tracked species formerly considered secure. Connell Memorial Herbarium, UNB, Fredericton NB. 2019. |
| 2 | Smith, M.E.M. 2008. AgCan Collection. Agriculture Canada, Charlottetown PE, 44 recs. |
| 2 | Thomas, A.W. 1996. A preliminary atlas of the butterflies of New Brunswick. New Brunswick Museum. |
| 2 | White, S. 2018. Notable species sightings, 2016-2017. East Coast Aquatics. |
| 1 | Amirault, D.L. 2000. Piping Plover Surveys, 1983-2000. Canadian Wildlife Service, Sackville, unpublished data. 70 recs. |
| 1 | Amirault, D.L. 2005. 2005 Peregrine Falcon Survey. Canadian Wildlife Service, Sackville, unpublished data. 27 recs. |
| 1 | Anon. Dataset of butterfly records for the Maritime provinces. Museum of Comparative Zoology, Harvard University. 2017. |
| 1 | Barney, T. 2020. Text message to Sean Blaney from Ted Barney with photograph of large Snapping Turtle at White Birch Impoundment, Westmorland Co., NB. pers. comm., 1 record. |
| 1 | Basquill, S. P. 2008. Nova Scotia Dept of Natural Resources. |
| 1 | Basquill, S.P. 2004. <i>C. americana</i> and <i>Sedum</i> sp records, 2002. Pers. comm. to C.S. Blaney. 2 recs, 2 recs. |
| 1 | Basquill, S.P. 2011. Field observations & specimen collections, 2010. Nova Scotia Department of Natural Resources, Pers. comm. , 8 Recs. |
| 1 | Basquill, S.P. 2012. 2012 Bryophyte specimen data. Nova Scotia Department of Natural Resources, 37 recs. |
| 1 | Basquill, S.P. 2018. Various specimens. NS DNR field work. NS Department of Natural Resources, 10. |
| 1 | Belliveau, A.G. E.C. Smith Herbarium Specimen Database 2019. E.C. Smith Herbarium, Acadia University. 2019. |
| 1 | Benedict, B. Connell Herbarium Specimen Database Download 2004. Connell Memorial Herbarium, University of New Brunswick. 2004. |
| 1 | Benedict, B. Connell Herbarium Specimens. Digital photos. University New Brunswick, Fredericton. 2005. |
| 1 | Benjamin, L.K. 2003. <i>Cypridium arietinum</i> in Cogmagun River NS. Pers. comm. to S. Blaney, 1 rec. |
| 1 | Blaney, C.S. 1999. Fieldwork 1999. Atlantic Canada Conservation Data Centre. Sackville NB, 292 recs. |
| 1 | Blaney, C.S. 2014. 2014 Bank Swallow colony observation, Westcock, NB. Atlantic Canada Conservation Data Centre. |
| 1 | Bonnyman, Vanessa. 2021. Eastern Wood-Pewee Observation on Bonshaw Trail, PEI. |
| 1 | Brach, A.R. 2019. Correspondence to Sean Blaney regarding <i>Calamagrostis cinnoides</i> specimen from Halifax NS. pers. comm., Harvard University Herbaria, 1 record. |
| 1 | Bruce, J. 2014. 2014 Wood Turtle email report, Nine Mile River, NS. NS Department of Natural Resources. |
| 1 | Clayden, S.R. 2006. <i>Pseudevernia cladonia</i> records. NB Museum. Pers. comm. to S. Blaney, Dec, 4 recs. |
| 1 | Clayden, S.R. 2020. Email to Sean Blaney regarding <i>Pilophorus cereus</i> and <i>P. fibula</i> at Fidele Lake area, Charlotte County, NB. pers. comm., 2 records. |
| 1 | COSEWIC (Committee on the Status of Wildlife in Canada). 2013. COSEWIC Assessment and Status Report on the Eastern Waterfan <i>Peltigera hydrothyria</i> in Canada. COSEWIC, 46 pp. |
| 1 | Crowell, A. 2004. <i>Cypridium arietinum</i> in Weir Brook, Hants Co. Pers. comm. to S. Blaney, 1 rec. |
| 1 | Curley, F.R. 2021. <i>Nymphalis l-album</i> record from near Belfast PEI. Pers. comm. to J. Klymko. |
| 1 | deGooyer, K. 2019. Snapping Turtle and Eastern White Cedar observations. Nova Scotia Environment. |
| 1 | Eastman, A. 2019. Snapping Turtle observation at Brookfield, Colchester Co. NS. Halifax Field Naturalists Nova Scotia Nature Archive Facebook Page, 1 record. |
| 1 | Edge, Thomas A. 1984. Status report on the Atlantic Whitefish (<i>Coregonus huntsmani</i>). Committee on the Status of Endangered Wildlife in Canada. |
| 1 | Elizabeth Spence. 2020. Email from Elizabeth Spence to John Klymko about the occurrence of a Wood Turtle in Westmorland County, New Brunswick. Pers. comm. |
| 1 | Gerriets, S.H. 1997-2001. Element Occurrence Database. Atlantic Canada Conservation Data Centre, Sackville NB, 1 rec. |
| 1 | Giberson, D. 2008. UPEI Insect Collection. University of Prince Edward Island, 157 recs. |
| 1 | Golder Associates Ltd. 2021. Black Ash location from Goff's Quarry Expansion Environment Assessment, 2017. Golder Associates Ltd., 1 record. |
| 1 | Goltz, J.P. 2001. Botany Ramblings April 29-June 30, 2001. N.B. <i>Naturalist</i> , 28 (2): 51-2. 8 recs. |
| 1 | Harris, Megan. 2018. Miscellaneous <i>Sorex palustris</i> record. Pers. comm. to S. Blaney. |
| 1 | Haughian, S.R. 2018. Description of <i>Fuscopannaria leucosticta</i> field work in 2017. New Brunswick Museum, 314 recs. |
| 1 | Hill, N.; Manning, I. 2020. Wild Leek observation, Cornwallis River, NS, floodplain. pers. comm. to J. Churchill. |
| 1 | Hill, N.M. 2013. email communications to Sean Blaney and David Mazerolle regarding the discovery of <i>Listera australis</i> populations at Black River Lake and Middlewood. , 2. |
| 1 | Jacques Whitford Ltd. 2003. Cananda Lily location. Pers. Comm. to S. Blaney. 2pp, 1 rec, 1 rec. |
| 1 | Kelly, Glen 2004. Botanical records from 2004 PEI Forestry fieldwork. Dept of Environment, Energy & Forestry, 71 recs. |
| 1 | Klymko, J. Henry Hensel's Butterfly Collection Database. Atlantic Canada Conservation Data Centre. 2016. |
| 1 | Klymko, J.J.D. 2012. Insect field work & submissions. Atlantic Canada Conservation Data Centre, 852 recs. |
| 1 | Lautenschlager, R.A. 2010. Miscellaneous observations reported to ACCDC (zoology). Pers. comm. from various persons, 2 recs. |
| 1 | Lautenschlager, R.A. 2020. Email to John Klymko about fisher sighting in Baie Verte. Klymko, J. (ed.) Lautenschlager, R.A., pers. comm. |
| 1 | Macaulay, M. 2008. Email to Sean Blaney regarding rich hardwood floodplain site at Howards Pool, Wallace River, NS. |

| # recs | CITATION |
|--------|--|
| 1 | MacAuley, M. 2020. Email to Sean Blaney regarding <i>Agalinis pauperula</i> var. <i>parviflora</i> at Malagash Station, NS. pers. comm., 2 records. |
| 1 | MacKinnon, D.; Wright, P.; Smith, D. 2014. 2014 Common Tern email report, Eastern Passage, NS. NS Department of Environment. |
| 1 | MacPhail, V. Bee and syrphid specimens from MSc research. Pers. comm., J. Klymko. 2006. |
| 1 | Majka, C.G. & McCorquodale, D.B. 2006. The Coccinellidae (Coleoptera) of the Maritime Provinces of Canada: new records, biogeographic notes, and conservation concerns. <i>Zootaxa</i> . <i>Zootaxa</i> , 1154: 49–68. 7 recs. |
| 1 | Mazerolle, D. 2003. Assessment of Seaside Pinweed (<i>Lechea maritima</i> var. <i>subcylindrica</i>) in Southeastern New Brunswick. Irving Eco-centre, la Dune du Bouctouche, 18 recs. |
| 1 | McKendry, Karen. 2016. Rare species observations, 2016. Nova Scotia Nature Trust, 19 recs. |
| 1 | Miller, D.G. 2013. Peregrine Falcon nesting information from birdingnewbrunswick.ca . birdingnewbrunswick.ca . |
| 1 | Morrison, Annie. 2010. NCC Properties Fieldwork: June-August 2010. Nature Conservancy Canada, 508 recs. |
| 1 | NatureServe Canada. 2018. iNaturalist Butterfly Data Export . iNaturalist.org and iNaturalist.ca . |
| 1 | Neily, P.D. Plant Specimens. Nova Scotia Dept Natural Resources, Truro. 2006. |
| 1 | Neily, T.H. & Pepper, C.; Toms, B. 2019. Boreal Felt Lichen Observation, April 2019. Mersey Tobeatic Research Institute. |
| 1 | Neily, T.H. 2004. <i>Hepatica nobilis</i> var. <i>obtusata</i> record for Falmouth NS. Pers. comm. to C.S. Blaney, 1 rec. |
| 1 | Neily, T.H. 2013. Email communication to Sean Blaney regarding <i>Agalinis pauperula</i> observations made in 2013 in Nova Scotia. , 1 rec. |
| 1 | Newell, R.E. 2004. <i>Hepatica nobilis</i> var. <i>obtusata</i> record. Pers. comm. to S. Blaney, 1 rec. |
| 1 | Niel, K. & Majka, C. 2008. New Records of Tiger Beetles (Coleoptera: Carabidae: Cicindelinae) in Nova Scotia. <i>Journal of the Acadian Entomological Society</i> , 4: 3-6. |
| 1 | Parker, M. 2018. East Coast Aquatics ACCDC 2018 Report. East Coast Aquatics, 12 records. |
| 1 | Payzant, P. 2018. Satyr Comma record from Bible Hill, NS. https://novascotiabutterflies.ca . |
| 1 | Popma, K. 2001. Phalarope & other bird observations in Westmorland Co. , Pers. comm. to K.A. Bredin. 5 recs. |
| 1 | Quigley, E.J. 2021. Email to Sean Blaney regarding Eastern White Cedar (<i>Thuja occidentalis</i>) stand near Shinimicas Bridge. NSDLF, 1 record. |
| 1 | Robinson, C.B. 1907. Early intervale flora of eastern Nova Scotia. <i>Transactions of the Nova Scotia Institute of Science</i> , 10:502-506. 1 rec. |
| 1 | Sabine, M. 2016. Black Ash records from NB DNR permanent forest sampling Plots. New Brunswick Department of Natural Resources, 39 recs. |
| 1 | Scott, F.W. 1988. Status Report on the Southern Flying Squirrel (<i>Glaucomys volans</i>) in Canada. Committee on the Status of Endangered Wildlife in Canada, 2 recs. |
| 1 | Skevington, Jeffrey H. 2020. Syrphid records used for the Field Guide to the Flower Flies of Northeastern North America. Canadian National Collection of Insects. |
| 1 | Sollows, M.C. 2008. NBM Science Collections databases: herpetiles. New Brunswick Museum, Saint John NB, download Jan. 2008, 8636 recs. |
| 1 | Speers, L. 2001. Butterflies of Canada database. Agriculture & Agri-Food Canada, Biological Resources Program, Ottawa, 190 recs. |
| 1 | Spicer, C.D. 2004. Specimens from CWS Herbarium, Mount Allison Herbarium Database. Mount Allison University, 5939 recs. |
| 1 | Toms, Brad. 2022. Non-Lichen Observations from Lichen SMP and NCC Property Searches. Mersey Tobeatic Research Institute. |
| 1 | Webster, R.P. 2004. Lepidopteran Records for National Wildlife Areas in New Brunswick. Webster, 1101 recs. |
| 1 | Whittam, R.M. 1999. Status Report on the Roseate Tern (update) in Canada. Committee on the Status of Endangered Wildlife in Canada, 36 recs. |
| 1 | Williams, M. Cape Breton University Digital Herbarium. Cape Breton University Digital Herbarium. 2013. |
| 1 | Wilson, G. 2013. 2013 Snapping Turtle email report, Wentworth, NS. Pers. comm. |
| 1 | Zahavich, J. 2017. Canada Warbler and Olive-sided Flycatcher records 2017. Island Nature Trust, 14 recs. |
| 1 | Zahavich, J. 2018. Canada Warbler and Olive-sided Flycatcher records 2018. Island Nature Trust, 14 recs. |