APPENDIX I

WILDLIFE PHOTO LOG



Photo 1: Placement of CAM1.



Photo 2: Viewpoint of CAM1.



Photo 3: Fisher observed by CAM1.



Photo 4: North American Deer Mouse observed by CAM1.



Photo 5: Red Squirrel observed by CAM1.



Photo 6: Coyote observed by CAM1.



Photo 7: Raccoon observed by CAM1.



Photo 8: Blue Jay observed by CAM1.



Photo 9: American Robin observed by CAM1.



Photo 10: Striped Skunk observed by CAM1.




Photo 11: American Black Bear observed by CAM1.



Photo 12: Placement of CAM2.



Photo 13: Viewpoint of CAM2.



Photo 14: North American Porcupine observed by CAM2.



Photo 15: White-tailed Deer observed by CAM2.



Photo 16: Coyote observed by CAM2.



Photo 17: American Black Bear observed by CAM2.



Photo 18: American Black Bear cub observed by CAM2.



Photo 19: Bobcat observed by CAM2.



Photo 20: Downstream section of Eel Weir Brook where Wood Turtle surveys were conducted.



Photo 21: Example of streamside foraging habitat surveyed for Wood Turtles along Eel Weir Brook.



Photo 22: Example of basking rocks observed during Wood Turtle surveys along Eel Weir Brook.



Photo 23: Example of streamside foraging habitat surveyed for Wood Turtles.



Photo 24: Example of deep over-wintering pools observed during Wood Turtles surveys along Eel Weir Brook.



Photo 25: Example of streamside foraging habitat observed during Wood Turtle surveys along Eel Weir Brook.



Photo 26: Example of deep over-wintering pools observed during Wood Turtles surveys in a large fen along Eel Weir Brook.



Photo 27: Example of amphibian eggs observed during Wood Turtles surveys along Eel Weir Brook.



Photos 28-31: Examples of locations selected for breeding bird surveys.



Photo 32: Placement and viewpoint of the Audiomoth bird acoustic monitor.



Photos 33-34: Placement and viewpoint of the Audiomoth bat acoustic monitor.







Photos 35-37: Examples of roadside habitat surveyed during PGI transects.







Photos 38-40: Examples of off-road habitat surveyed during PGI transects.



Photo 41: Example of deer bones found during PGI transects.



Photo 42: Deer jawbone observed during PGI transects.



Photo 43: Example of Coyote scat observed during PGI transects.



Photo 44: Examples of Snowshoe Hare and White-tailed Deer pellets observed during PGI transects.



Photo 45: Example of North American Porcupine scat observed during PGI transects.



Photo 46: Example of White-tailed Deer tracks observed during PGI transects.



Photo 47: Examples of White-tailed Deer browsing activity observed during PGI transects.



Photos 48-49: Example of Beaver gnawing activity and Pileated Woodpecker feeding cavities observed during PGI transects.



Photo 50: Example of small mammal tracks observed during Mainland Moose winter tracking surveys.



Photo 51: Coyote tracks and evidence of prey dragging observed during Mainland Moose winter tracking surveys.



Photo 52: Coyote tracks observed during Mainland Moose winter tracking surveys.



Photo 53: White-tailed Deer tracks observed crossing WC1 during Mainland Moose winter tracking surveys.



Photo 54: Snowshoe Hare tracks observed during Mainland Moose winter tracking surveys.



Photo 55: White-tailed Deer remains observed during Mainland Moose winter tracking surveys.



Photo 56: Assorted tracks observed during Mainland Moose winter tracking surveys.



Photo 57: Coyote scat observed during Mainland Moose winter tracking surveys.



Photo 58: Likely predation event showing small mammal tracks leading to talon marks.